

DEPARTMENT OF COMMERCE

CIRCULAR

OF THE

BUREAU OF STANDARDS

S. W. STRATTON, DIRECTOR

No. 81

BIBLIOGRAPHY OF SCIENTIFIC LITERATURE RELATING TO HELIUM

[Second edition]
DECEMBER 21, 1922



PRICE, 10 CENTS

Sold only by the Superintendent of Documents, Government Printing Office
Washington, D. C.

WASHINGTON
GOVERNMENT PRINTING OFFICE
1923

DEPARTMENT OF COMMERCE

CIRCULAR
OF THE
BUREAU OF STANDARDS

S. W. STRATTON, DIRECTOR

No. 81

**BIBLIOGRAPHY OF SCIENTIFIC LITERATURE
RELATING TO HELIUM**

[Second edition]
DECEMBER 21, 1922



PRICE, 10 CENTS

Sold only by the Superintendent of Documents, Government Printing Office
Washington, D. C.

WASHINGTON
GOVERNMENT PRINTING OFFICE
1923

BIBLIOGRAPHY OF SCIENTIFIC LITERATURE RELATING TO HELIUM.¹

ABSTRACT.

A guide to the useful literature relating to helium from the time of its discovery to January 1, 1922.

CONTENTS.

	Page.
I. Introduction.....	4
II. Arrangement of titles.....	5
III. Discovery and identification.....	5
1. Discovery.....	5
2. Elementary nature of helium.....	6
IV. Occurrence of helium.....	6
1. In minerals.....	6
2. Occurrence in mineral waters.....	8
3. Occurrence in natural gas.....	9
4. Occurrence in air.....	10
V. Formation of helium.....	10
1. From radioactive substances.....	10
2. Supposed formation of helium from nonradioactive substances...	12
VI. Separation and purification of helium.....	12
VII. Analysis of helium.....	14
VIII. Properties of helium.....	14
1. Electrical and magnetic properties.....	14
2. Spectrum of helium.....	16
3. Optical properties of helium.....	20
4. Solubility and absorption of helium.....	21
5. Density and atomic weight of helium.....	22
6. Pressure-volume relations for gaseous helium.....	23
7. Diffusion and effusion of helium.....	24
8. Viscosity of helium.....	24
9. Specific heats and other thermal properties of helium.....	25
10. Thermal conductivity of helium.....	25
IX. Liquefaction and properties of liquid helium.....	26
X. Applications of helium.....	27
1. Application of helium to thermometry.....	27
2. Applications of liquid helium especially to determination of electrical properties of metals at low temperatures.....	27
3. Applications of helium to photometry, etc.....	28
4. Application of helium as a balloon gas.....	28
XI. Relation of helium to subatomic phenomena.....	29
XII. Miscellaneous and general.....	32

¹ Prepared by E. R. Weaver, Bureau of Standards, Washington, D. C.

I. INTRODUCTION.

Helium has probably been the most interesting of all the elements to the theoretical scientist on account of the romantic history of its discovery, its occurrence in a remarkable condition of solid solution in many minerals, its formation as a product of the disintegration of the radioactive elements, its liquefaction after a decade of unsuccessful attempts by some of the world's greatest experimenters, the attainment by its use of temperatures below those at which the resistances of pure metals vanish, its many unique physical properties, and the many important theoretical conclusions which have been drawn from its behavior.

All of these points of interest have been the subjects of very thorough investigation. The important developments of the future will probably be along the line of the applications of helium, many of which have already been suggested; but in order to make the most of these possible applications it is necessary to know the properties on which they are based. It is as a guide to these properties that, it is hoped, this bibliography will find its chief usefulness.

This bibliography was first prepared at the beginning of the development of helium for balloon-gas purposes and was intended as an aid in that enterprise. The first edition was issued in 1919 and was intended to cover the literature prior to January 1 of that year. This second edition is intended to cover the literature to January 1, 1922, and includes some references of later date. After its publication the first edition was found to be less complete than had been thought, and no claim is made for completeness in this second edition. Some articles have been purposely omitted. These include a small number of articles which, for one reason or another, have been considered of no importance or positively misleading, and reviews and other articles containing no original work which were published in inaccessible foreign journals and contained no material which was not available in English or American publications. Other material, especially much of that dealing directly with radioactivity, the alpha-particle, and the structure of the atom, has been omitted because it is only indirectly concerned with helium. It is difficult to make a sharp distinction between articles of this class which pertain to helium and those which do not. An effort has been made to include enough representative articles on these subjects to at least serve as an introduction to their study.

II. ARRANGEMENT OF TITLES.

The arrangement of material under each subhead has, in general, been such that closely related articles occur together in their chronological order. The bibliography is thus, in effect, a brief outline history of the subject. An exception to this arrangement is made in the case of articles on the occurrence of helium, which have been arranged alphabetically according to the authors' names. This was done because the papers by different authors are usually but slightly related to each other, and the chronological development seems of less importance than the bringing together of the papers, often numerous, of each author.

III. DISCOVERY AND IDENTIFICATION.

1. DISCOVERY.

- JANSSEN. *Compt. rend.*, **67**, p. 838; 1868: Discovery of new spectrum lines in sun, since found to belong to helium.
- FRANKLAND and LOCKYER. *Proc. Roy. Soc., Lond.*, **17**, p. 91; 1868: Announce the existence of an element in the sun unknown on earth and name it helium.
- PALMIERI. *Gazz. Chim. Ital.*, **12**, p. 556; 1882: Discovery of helium spectrum in rocks from Vesuvius.
- NASINI and ANDERLINI. *Atti. Accad. Lincei. Roma.* (5), **13**, I, p. 368; 1904: Recognition of helium in Vesuvius lavas confirming the discovery of Palmieri.
- HILLEBRAND, W. F. *Am. J. Sci.* (3), **40**, p. 384; 1890: Observed the presence in uraninite of gases since found to be helium and nitrogen. Condensed form of following reference.
- HILLEBRAND. *Bull. U. S. Geol. Survey*, No. 78, p. 43; 1890: Chemical and spectroscopic tests of gases obtained from uraninite led to conclusion that gas was nitrogen.
- HILLEBRAND. *Am. Jour. Sci.* (3), **42**, p. 390; 1891: New analyses of uraninite.
- RAMSAY. *Proc. Roy. Soc., Lond.*, **58**, pp. 65, 81; 1895: Reports partial identification of helium spectrum in gas from cleveite.
- LOCKYER. *Proc. Roy. Soc., Lond.*, **58**, p. 67; 1895: Partial confirmation of Ramsay's identification of helium.
- RAMSAY. *J. Chem. Soc.* 67, pp. 684, 1107; *Chem. News*, **71**, p. 151; 1895: Discovery of helium in cleveite.
- CROOKES. *Chem. News*, **71**, p. 151; 1895: Measured the wave lengths of the spectrum lines of the gas isolated from cleveite by Ramsay and identified the gas as helium.
- RUNGE and PASCHEN. *Chem. Ztg.*, **19**, p. 997; 1895: Observed double lines in spectrum of helium from minerals and cast doubt upon identity of mineral and solar helium.
- HUGGINS, W. *Chem. News*, **71**, p. 283; 1895: Differences in spectra of helium from minerals and in the sun.
- HUGGINS, W. *Chem. News*, **72**, p. 27; 1895: Observed double helium lines in solar spectrum.
- RUNGE and PASCHEN. *Math. natw. Mitt. Berlin*, p. 323; 1895: Identity of spectra of solar and mineral helium.
- WILDE. *Phil. Mag.* (5), **40**, p. 466; 1895: Identity of spectra of solar and mineral helium.
- LOCKYER. *Proc. Roy. Soc., Lond.*, **62**, p. 52; 1897: Existence of helium in fixed stars and nebulae.

- RAMSAY and TRAVERS. *Compt. rend.*, **126**, p. 1762; 1898: New atmospheric gases.
 RAMSAY. *Ann. chim. phys.* (7), **13**, p. 433; 1898: Discovery of helium.
 RAMSAY, W. *The Gases of the Atmosphere, the History of their discovery.* Book.
 London, Macmillan & Co. *Chem. Abs.*, **10**, p. 313; 1916.
 ABBOTT, C. G. *Smithsonian Report for 1918*, pp. 121-126; *Smithsonian publ. No.*
 2550; 1920: The discovery of helium and what came of it.
 LOCKYER, W. J. S. *Nature*, **105**, pp. 360-363; 1920: Helium, its discovery and
 applications.
 MOORE, R. B. *J. Frank. Inst.*, **191**, pp. 145-197; 1921: Helium, its history, prop-
 erties, and commercial development.

2. ELEMENTARY NATURE OF HELIUM.

[See also Formation of helium and Relation to subatomic phenomena.]

- RAMSAY. *Proc. Roy. Soc., Lond.* **60**, p. 53; 1896: Experiments which show the
 inactivity of helium and argon.
 RAMSAY. *Ber. d. chem. Ges.*, **31**, p. 3111; 1898: The newly discovered gases and
 their relation to the periodic law.
 RAMSAY and COLLIE. *Compt. rend.*, **123**, p. 214; *Nature*, **54**, p. 546; *Proc. Roy.*
Soc., Lond., **60**, p. 206; 1896: Homogeneity of argon and helium.
 RAMSAY and TRAVERS. *Proc. Roy. Soc., Lond.*, **62**, p. 316; 1897: Dispels the doubt
 as to the homogeneity of helium suggested in preceding reference.
 BRAUNER, B. *Chem. News*, **71**, p. 271; 1895.
 BRAUNER, B. *Chem. News*, **74**, p. 223; 1896.
 HILL, E. A. *Am. J. Sci.*, (3), **50**, p. 359; 1895.
 RAYLEIGH. *Chem. News*, **74**, p. 260; 1896.
 MATHEWS, A. P. *J. Phys. Chem.*, **17**, pp. 337-343; 1913; *Chem. Abs.*, **7**, 1899: Val-
 ence of the argon group as determined from molecular cohesion.
 SKAUPY, F. *Zeit. Physik.*, **3**, pp. 408-411; 1920; *Chem. Abs.*, **15**, p. 2024: Chemical
 affinity of the noble gases.
 ASTON, F. W. *Nature*, **105**, p. 8; 1920; *Chem. Abs.*, **14**, p. 1629: The constitution
 of the elements. Determination of the atomic weight of helium by the positive
 ray method.
 ASTON, F. W. *Sci. Progress*, **15**, pp. 212-222; 1920; *Chem. Abs.*, **15**, p. 335: Mass
 spectra and the atomic weights of the elements.
 BRINER, E. *J. Chim. Phys.*, **19**, pp. 9-10; 1921; *Chem. Abs.*, **16**, p. 181: Some nega-
 tive attempts at transformation of elements. Attempt to decompose iodine,
 hydrogen, and helium.

IV. OCCURRENCE OF HELIUM.²

1. IN MINERALS.

[See also Formation from radioactive substances.]

- ADAMS, E. P. *Am. J. Sci.* (4), **19**, p. 321; 1905: Helium in carnotite.
 BORDAS, F. *Compt. rend.*, **146**, pp. 628-630, 1908; *Chem. Abs.*, **2**, p. 1926: Search
 for small quantities of helium in minerals.
 BORDAS. *Compt. rend.*, **146**, pp. 896-898; 1908; *Chem. Abs.*, **2**, p. 2043: Helium in
 minerals containing uranium.
 BROGGER, W. C. *Pharmacia*, **1**, pp. 49-53, 65-70; 1904: Ores of uranium containing
 helium and radium.
 CLEVE. *Compt. rend.*, **120**, p. 834, 1895; *Chem. News*, **71**, p. 201, 1895: Occurrence
 of helium in cleveite.

² The material under this heading is arranged alphabetically according to authors' names because of the general slight relation between papers by different authors.

- COLLIE, J. N., and TRAVERS, M. W. J. Chem. Soc., **67**, p. 684; 1895: Helium a constituent of certain minerals.
- DEBIERNE, A. Ann. Physik (9), **2**, pp. 478-488; 1904: Preparation of helium from fluorspar crystals. Contains also an account of unsuccessful efforts to obtain helium from other sources.
- KITCHIN, E. S., and WINTERSON, W. G. J. Chem. Soc., **89**, pp. 1568-1575; 1906: Malacone, a silicate of zirconium containing helium.
- KOHLSCHUTTER, V. Ann. Chem., **317**, pp. 158-189; 1901: Helium in uranium minerals.
- KOHLSCHUTTER and VOGDT. Ber. d. chem. Ges., **38**, pp. 1419-1430, 2992-3002; 1905: Helium in uranium minerals.
- LANGÉ, H. Zeit. Naturw., **82**, pp. 1-34, 1910; Chem. Abs., **4**, p. 2920: Studien über die Zusammensetzung heliumführender Mineralien. (Studies on the structure of helium-bearing minerals.) Issued as separate publication.
- LANGLET. Z. anorg. Chem., **10**, p. 289; 1895: Helium in cleveite.
- LOCKYER. Compt. rend., **120**, p. 1103; 1895; Chem. News, **72**, p. 283; 1895: Occurrence of helium in minerals.
- MAGLI, G. Rend. soc. chim. ital. (2), **5**, pp. 420-423; 1914; Chem. Abs., **9**, p. 1005: Helium in titanite. Occurs in amounts proportional to the radioactivity of the mineral.
- MOISSAN and DESLANDERS. Compt. rend., **126**, p. 1689; 1898: Helium in cerite.
- MOSS. Trans. Roy. Dublin Soc. (2), **8**, p. 153; 1904: Removal of helium from minerals by evacuation. Can obtain only a little over 1 per cent of total by this method.
- VON OEFELE, F. Pharm. Zentralhalle, **57**, pp. 83-84; 1916; J. Chem. Soc., **110**, II, p. 284; 1916; Chem. Abs., **10**, p. 2654: Helium in samarskite and its relation to other elements present.
- PIUTTI, A. Radium, **7**, pp. 146-149; 1910; Chem. Abs., **4**, p. 3055: Nonradioactive minerals containing helium.
- PIUTTI. Radium, **7**, pp. 178-179; 1910; Chem. Abs., **5**, p. 247: Helium in recent minerals.
- PIUTTI. Nature, **84**, pp. 543-544; 1910; Chem. Abs., **5**, p. 626: The absorption of helium in salts and minerals. A hypothetical explanation of the accumulation of helium in certain minerals.
- PIUTTI. Radium, **8**, pp. 204-205; 1911; Chem. Abs., **6**, p. 1255: The presence of helium in autinite and the period of life of ionium.
- PIUTTI. Radium, **10**, pp. 165-168; 1913; Chem. Abs., **7**, p. 3076: Helium in glaucinum minerals.
- PIUTTI. Atti Acad. Lincei, **22**, pp. 140-144; 1913; Chem. Abs., **7**, p. 2352: Helium and beryllium minerals.
- RAMSAY, W. Proc. Roy. Soc., Lond., **59**, p. 325; 1895: Helium, a gaseous constituent of certain minerals.
- RAMSAY. Comp. rend., **120**, p. 1049; 1895; Ann. chim. phys. (7), **13**, p. 433; 1898: Helium in meteorites.
- RAMSAY and TRAVERS. Proc. Roy. Soc., Lond., **60**, p. 442; 1896: Occurrence of helium in minerals.
- STRUTT, R. J. Nature, **75**, p. 271; 1907; Chem. Abs., **1**, p. 817: Helium and argon in common rocks.
- STRUTT. Nature, **75**, p. 390; 1907: An occurrence of helium in the absence of radio activity.
- STRUTT. Proc. Roy. Soc., Lond., **76**, pp. 80-101; 1908: Helium and argon in rocks.
- STRUTT. Proc. Roy. Soc., Lond., **80**, pp. 56-57; 1908; Chem. Abs., **2**, p. 944, Radium, **5**, pp. 202-211; 1908: The association of helium and thorium in minerals.

- STRUTT. *Proc. Roy. Soc., Lond.*, **80**, pp. 572-594; 1908; *Chem. Abs.*, **3**, p. 2651: Helium and radioactivity in minerals.
- STRUTT. *Proc. Roy. Soc., Lond.*, **81**, pp. 278-279; 1908; *Chem. Abs.*, **3**, p. 2651: Helium in saline minerals and its probable connection with potassium.
- STRUTT, R. J. *Proc. Roy. Soc., Lond.*, **81**, pp. 272-277; 1908; *Chem. Abs.*, **3**, p. 615: The accumulation of helium in geological time.
- STRUTT. *Chem. News*, **99**, pp. 145-146; 1909; *Chem. Abs.*, **3**, p. 2531: The leakage, of helium from radioactive minerals.
- STRUTT. *Proc. Roy. Soc., Lond.*, **83**, pp. 76-99, 298-301; 1910; *Chem. Abs.*, **4**, pp. 1146, 1424: The accumulation of helium in geological time.
- STRUTT. *Proc. Roy. Soc., Lond.*, **84**, pp. 194-196; 1910; *Chem. Abs.*, **5**, p. 28: The accumulation of helium in geological time.
- STRUTT. *Nature*, **85**, p. 6; 1911; *Chem. Abs.*, **5**, p. 626: Helium and geological time. An answer to Piutti's hypothesis of absorption to account for the presence of helium in rocks.
- THOMSON, J. *Zeit. Physik. Chem.*, **25**, p. 112; 1898: Occurrence of helium in minerals. Chiefly calcium fluoride.
- TRAVERS. *Nature*, **71**, p. 248; 1905: Helium occurs in radioactive minerals in form of supersaturated solid solution.
- TSCHERNIK, J. *J. Russ. Phys. Chem. Soc.*, **29**, p. 291; 1899: Helium in cerium-bearing minerals from the Caucasus.
- VALENTINER, S. *Kali*, **6**, pp. 1-3; *Chem. Abs.*, **8**, p. 480; *Neues Jahrb. Min. Geol.*, 1913, I Ref., p. 195: Helium content of blue rock salt.
- WELLS, R. C. J. *Frank. Inst.*, **189**, pp. 779-780; 1920; *Chem. Abs.*, **14**, p. 2456: Note on brannerite, a uranium mineral containing helium.
- WHERRY, E. T. *Am. Mineral*, **2**, pp. 105-108; 1917; *Chem. Abs.*, **11**, p. 2570: The occurrence of the native elements. (Including helium.)
- WILDE, H. *Phil. Mag.* (5), **40**, p. 466; 1895: The occurrence of helium in cleveite.

2. OCCURRENCE IN MINERAL WATERS.

- BOUCHARD and TROOST. *Compt. rend.*, **121**, p. 392; 1895: Helium in a spring at Cauteret in the Pyrenees.
- CZAKO, E., and LAUTENSCHLAGER. *Chem. News*, **108**, p. 16; 1913; *Chem. Ztg.*, **37**, p. 936; 1913; *Chem. Abs.*, **7**, p. 3450: Helium content of gases from Hôt Spring, at Wildbad, in the Black Forest. Gases contained 0.71 per cent helium.
- DEWAR, J. The discovery of helium and radium in the hot springs of Bath, England, 14 pp. Bath Herald Office. 1904.
- EWERS. *Physik. Zeit.*, **7**, p. 224; 1906: Helium and argon in hot springs.
- HERRMANN, A., and PESENDORFER, F. *Physik. Zeit.*, **6**, pp. 70-71; 1905: Radioactivity of the gases evolved from the Karlsbad springs.
- KAYSER, H. *Chem. News*, **72**, p. 223; 1895; *Chem. Ztg.*, **19**, p. 1549; 1895: Helium in the gases from springs at Wildbad, in the Black Forest.
- MOUREU, C. *Compt. rend.*, **121**, p. 819; 1895: Helium in spring water at Maizieres.
- MOUREU. *Compt. rend.*, **135**, p. 1335; 1902: Helium in the spring Vieille Quelle, in the Pyrenees.
- MOUREU. *Compt. rend.*, **139**, pp. 852-855; 1904: The chemical composition of the mixture of radioactive gases liberated from the waters of certain hot springs.
- MOUREU. *Compt. rend.*, **142**, p. 1155; 1906.
- MOUREU and BICQUARD. *Compt. rend.*, **143**, pp. 795-797; 1906: Helium in certain springs. Gas from one of the springs contained 5.34 per cent helium.
- MOUREU and BICQUARD. *Compt. rend.*, **146**, pp. 435-437; 1908: Recent researches on the rare gases from hot springs.
- MOUREU. *Bull. Soc. Chim.* (4), **9**, pp. 1-25; 1911; *Rev. gen. sci.* **49**, pp. 65-76, 1911: Investigations on the rare gases from hot springs.

- NASINI, R. *Atti Acad. Lincei. Roma* (5), **13**, I, pp. 217, 367; 1904; Helium in certain Italian springs.
- PESENDORFER, F. *Chem. Ztg.*, **29**, p. 359; 1905: Helium in the Karlsbad springs.
- PRYTZ and THORKELSSON. *Kgl. Danske. Vidensk. Selsk. Forh.*, p. 317; 1905: The occurrence of helium in varna island springs.
- RAYLEIGH. *Chem. News*, **72**, p. 223; 1895: Helium in gases from the springs at Bath.
- SIEVEKING and LAUTENSCHLAGER. *Physik. Zeit.*, **13**, pp. 1043-1051; 1912; *Chem. Abs.*, **7**, 1841; *Ber. physik. Ges.*, **14**, p. 910; 1913; Helium in hot springs and natural gases. No direct relation between radioactivity and helium content was found.
- TROOST and OUVARD. *Compt. rend.*, **121**, p. 798; 1895; Helium in spring water at Cauterets.

3. OCCURRENCE IN NATURAL GAS.

- CADY, H. P., and MCFARLAND, D. F. *Trans. Kan. Acad. Sci.*, **26**, Part II, p. 802; 1907; *Sci.*, **24**, p. 344; 1906; *Chem. Abs.*, **1**, p. 1528; Helium in Kansas natural gas. Some samples contained as much as 2 per cent. Geological zones of approximately equal helium content were located.
- CADY, H. P., and MCFARLAND, D. F. *J. Am. Chem. Soc.*, **29**, pp. 1524-1536; 1907; *Chem. Abs.*, **2**, p. 386: The occurrence of helium in natural gas and the composition of natural gas. Helium is found in practically all natural gas in amounts which increase, in general, with increasing nitrogen and decreasing hydrocarbon content.
- CZAKO, E. *Zeit. anorg. Chem.*, **82**, pp. 249-277; 1913; *Chem. Abs.*, **7**, p. 3450: The helium content and radioactivity of natural gases. The production of helium discharged yearly from two of the wells examined would require the disintegration of 165,000 and 28,000 tons of Ra, respectively.
- ERDMANN, E. *Ber. d. chem. Ges.*, **43**, pp. 777-782; 1910; *Chem. Abs.*, **4**, p. 1957: Helium-containing gases of the German potash beds.
- HARRIS, J. E. G. *J. Inst. Petroleum Tech.*, **7**, No. 27, pp. 14-16; 1921: Helium in a sample of natural gas from Egypt.
- MCLENNAN, J. C. *Can. Dept. of Mines Bull.*, **31**, 72 pp.; 1920: Report on sources of helium in the British Empire.
- MILNER, H. B. *Nature*, **109**, pp. 112; 1922; *Chem. Abs.*, **16**, p. 1132: Helium in natural gas.
- MOUREAU, C., and LÉPAPE, A. *Compt. rend.*, **155**, pp. 197-200; 1912; *Chem. Abs.*, **6**, p. 3075: Some natural gaseous mixtures particularly rich in helium. Discusses gases from eight French sources.
- MOUREAU, C., and LÉPAPE, A. *Compt. rend.*, **158**, pp. 598-603; 1913; *Chem. Abs.*, **8**, p. —; 1699: Helium from fire damp and the radioactivity of coal. Helium evolved from one mine is equal to 12 m.³ per day. Radioactivity of gas and coal does not account for it.
- ROGERS, G. S. *U. S. Geol. Survey Professional Paper No. 121*, 110 pp.; 1921: Helium-bearing natural gas.
- SATTERLY, J., and MCLENNAN, J. C. *Trans. Roy. Soc., Can.*, **12**, III pp. 153-160; 1918-1919; *Chem. Abs.*, **13**, p. 2309: The radioactivity of the natural gases of Canada.
- SEIBEL, C. W. Paper read before meeting of the Am. Chem. Soc., Apr. 10-14, 1917; *Abs. in Met. Chem. Eng.*, May, 1917: Helium and associated elements in Kansas natural gas.
- VOLLER, A., and WALTER, B. *Hamburger Wiss. Inst.*, **28**, p. —; 1910; *Chem. Abs.*, **5**, p. 3510; *Petroleum*, **6**, p. 1062: Helium and argon in the natural gas of Neuengamme. Contained 0.01-0.02 per cent helium.

4. OCCURRENCE IN AIR.

- RAYLEIGH and RAMSAY. *Proc. Roy. Soc., Lond.*, **59**, p. 198; 1896; *Proc. Roy. Soc., Lond.*, **60**, p. 206; 1896: Helium in the air. Negative results.
- TRAVERS. *Proc. Roy. Soc., Lond.*, **60**, p. 449; 1896: Separation of helium from the air.
- KAYSER, H. *Chem. Ztg.*, **19**, p. 1549; 1895: Helium in the air.
- FRIEDLANDER. *Z. physik. Chem.*, **19**, p. 657; 1896: Helium in the air. Estimated to be one part per billion.
- CROOKES, W. *Chem. News*, **78**, p. 198; 1898: Helium in the air.
- RAMSAY and TRAVERS. *Proc. Roy. Soc., Lond.*, **67**, p. 329; 1900: Helium in the air. Isolated by fractionation.
- DEWAR, J. *Proc. Roy. Soc., Lond.*, **68**, pp. 360-366; 1901; *Proc. Roy. Soc., Lond.*, **74**, 127; 1904: Separation of helium from the air.
- RAMSAY. *Proc. Roy. Soc., Lond.*, **76**, p. 111; 1905: Amount of helium in the air. One part in 245,300 by volume.
- BORDAS, F., and TOUPLAIN. *Compt. rend.*, **147**; pp. 591-594; 1908: Helium in the air.
- NASINI, ANDERLINI, and SALVADORI. *Mem. Accad. Lincei* (5), **5**, pp. 25-82; 1905: Gases from Vesuvius and Campo Flegrei.
- PIUTTI, A. Radium, **7**, pp. 142-146; 1910; *Chem. Abs.*, **5**, p. 247: Helium in the air of Naples and Vesuvius.

V. FORMATION OF HELIUM.

1. FROM RADIOACTIVE SUBSTANCES.

- RAMSAY and SODDY. *Proc. Roy. Soc., Lond.*, **72**, p. 204; 1903; *Zeit. phys. Chem.* **47**, pp. 490-494; 1904; *Physik Zeit.*, **4**, p. 229; 1903: The production of helium by radium.
- RAMSAY and SODDY. *Proc. Roy. Soc., Lond.*, **73**, pp. 346-358; 1904; *Zeit. physik. Chem.*, **48**, pp. 682-696; 1904; *Physik. Zeit.* **5**, pp. 349-356; 1904: Experiments on the production of helium from radium.
- STARK, J. *Natur. Rundschau*, **18**, pp. 429-430; 1903: The formation of helium from radium.
- STARK, J. Separate publication 1903: Dissozierung und Umwandlung chemischer, Atome. (The dissociation and transformation of chemical atoms.)
- DEWAR and CURIE. *Compt. rend.*, **138**, p. 190; 1904; *Chem. News*, **89**, p. 85; 1904: Formation of helium from radium.
- HIMSTEDT and MEYER. *Ann. Physik.* (4), **15**, p. 184; 1904: Formation of helium from radium.
- JENDRIKSON. *Physik. Zeit.*, **5**, p. 214; 1904: Formation of helium from radium.
- SCHENK. *Sitz. Ber. Akad. Wiss., Berlin*, **37**; 1904: Formation of helium from radium.
- RUTHEFORD. *Phil. Mag.* (6), **10**, p. 290; 1905; *Arch. sc. phys. nat.*, **19**, pp. 31, 125; 1905: Formation of helium from radium.
- DEBIERNE. *Compt. rend.*, **141**, p. 383; 1905: Helium from radium salts and actinium.
- GIEZEL, F. *Ber. d. chem. Ges.*, **38**, p. 2299; 1905: Proof of formation of helium from radium bromide.
- FREUND, M. *Jahr. physik. Ver.* 1904-1905, pp. 38-39; 1906: The transformation of radium into helium.
- CROOKES. *Chem. News*, **94**, p. 144; 1906: The production of helium from radium.
- DEWAR, J. *Proc. Roy. Soc., Lond.*, **81**, pp. 280-286; 1908; *Chem. News*, **98**, pp. 188-190; *Chem. Abs.*, **3**, pp. 281, 616: The rate of production of helium from radium.
- DEWAR, J. *Proc. Roy. Soc., Lond.*, **83**, p. 404; 1910; *Chem. Abs.*, **4**, p. 2410: Long-period determination of the rate of production of helium from radium. Found 0.463 mm.³/g. per day.

- RUTHERFORD, E., and BOLTWOOD, B. B. *Proc. Manch. Lit. Phil. Soc.* (6), **54**; 1910; *Chem. Abs.*, **4**, p. 1266: Production of helium by radium.
- BOLTWOOD and RUTHERFORD. *Phil. Mag.*, **22**, pp. 586-604; 1911; *Chem. Abs.*, **6**, p. 26: Production of helium from radium. Find rate of 156 cu. mm. per year per gram of radium.
- KAUFFMANN, H. *Zeit. angew. Chem.*, **17**, p. 1393; 1904: Formation of helium from radium emanation.
- HEMSTEDT and MEYER. *Ann. Physik* (4), **17**, pp. 1005-1008; 1905: Formation of helium from radium emanation.
- HEMSTEDT and MEYER. *Ber. natf. Ges.*, **16**, pp. 10-12; 1906: Formation of helium from radium emanation.
- BRUNER, L., and BEKIER, E. *Physik Z.*, **15**, pp. 240-241; 1914; *Chem. Abs.*, **8**, p. 1700: Attempt to reverse the reaction $\text{RaEm} = \text{RaA} + \text{He}$ ion by means of an electric discharge in helium gas.
- MARCKWALD, W. *Physik. Z.*, **15**, pp. 440-441; 1914; *Chem. Abs.*, **8**, p. 2845: Experiments on the decomposition of radium emanation in a helium atmosphere. No indication that disintegration of radium emanation can be retarded was found.
- RAMSAY, W. *Z. angew. Chem.*, **24**, pp. 1304-1307; 1908; *Chem. Abs.*, **2**, p. 2645: The radioactive gases and their relation to the noble gases of the atmosphere.
- RAMSAY, W. *Arch. sci. phys. nat.*, **9**, pp. 237-262; 1909: The inert gases of the atmosphere and their derivation from the emanations of the radioactive substances.
- GIESEL, F. *Ber. d. chem. Ges.*, **40**, p. 3011-3017; 1907; *Chem. Abs.*, **2**, 755: The first decomposition products of actinium and the formation of helium from actinium.
- BOLTWOOD, B. B. *Proc. Roy. Soc., Lond.*, **85**, pp. 77-81; 1910; *Chem. Abs.*, **5**, p. 3539: Report on the separation of ionium and actinium from certain residues and on the production of helium by ionium.
- SODDY, F. *Nature*, **79**, p. 129; *Radium*, **5**, pp. 361-362; 1908: Production of helium from uranium.
- SODDY. *Umschau*, **13**, pp. 375-377; 1907: Production of helium from uranium and thorium.
- STRUTT, R. J. *Nature*, **81**, p. 158; 1909; *Chem. Abs.*, **4**, p. 412: Rate of helium production from the complete series of uranium products. Finds 10.4×10^{-8} cm.³ per year per gram of U_3O_8 .
- STRUTT, R. J. *Proc. Roy. Soc., Lond.*, **84**, pp. 379-388; 1910; *Chem. Abs.*, **5**, p. 627: Measurements of the rate of production of helium in thorianite and pitchblende.
- SODDY. *Phil. Mag.* (6), **16**, pp. 513-30; 1908; *Chem. Abs.*, **3**, p. 281: Attempts to detect the production of helium from the primary radio-elements. One result gave 2×10^{-12} gram helium per year per gram of thorium.
- RUTHERFORD, E. *Trans. Roy. Soc., Lond.*, **204**, pp. 169-217; 1904: The succession of changes in radioactive bodies.
- MEYER, G. *Zeit. Electrochem.*, **13**, pp. 375-377; 1907: Evolution of helium from radioactive substances.
- BOLTWOOD, B. B. *Am. J. Sci.* (4), **23**, pp. 77-78; 1907; *Physik. Zeit.*, **8**, pp. 97-141; 1907: The ultimate disintegration products of the radioactive elements.
- DEBIERNE, A. *Ann. Physik*, **2**, pp. 428-488; 1914; *Chem. Abs.*, **9**, p. 1428: Review of experiments on the production of helium by radioactive substances.
- HESS, F., and LAWSON, R. W. *Arch. sci. phys. nat.*, **46**, pp. 330-331; 1918; *Chem. Abs.*, **13**, p. 3069: Number of alpha particles emitted by radium.
- KELLER, H. F. *J. Frank. Inst.*, **116**, p. 213; 1908: The transformations of the elements. A popular account of radioactivity and related phenomena.
- RUTHERFORD, E. *Radioactive substances and their radiation*. 1913. Cambridge Univ. Press.

2. SUPPOSED FORMATION OF HELIUM FROM NONRADIOACTIVE SUBSTANCES.

- RAMSAY, W. J. Chem. Soc., **103**, pp. 264-266; 1913; Chem. Abs., **7**, p. 1441: The presence of helium in the gas from the interior of an X-ray bulb.
- COLLIE, J. N., and PATTERSON, H. Nature, **90**, pp. 653-654; 1913; Chem. Abs., **7**, p. 1441: Origin of helium and neon in vacuum tubes.
- THOMSON, J. J. Nature, **90**, pp. 645-647; 1913; Chem. Abs., **7**, p. 1441; Science, **37**, pp. 360-364; Mon., Sci. (5), **3**, pp. 628-629; 1913: The appearance of helium and neon in vacuum tubes. Explained as liberation from substances not radioactive unless aided by bombardment by cathode rays.
- STRUTT, R. J. Proc. Roy. Soc., Lond. **89**, pp. 499-506; 1914; Chem. Abs., **8**, p. 1236: Unsuccessful attempts to observe the production of neon or helium by electric discharge.
- COLLIE, PATTERSON, and MASSON. Proc. Roy. Soc., Lond. **91**, pp. 30-45; 1915; Chem. Abs., **9**, p. 20: The production of helium and neon by the electric discharge. Argument to prove production.
- DEBIERNE, A. Ann. Physik., **2**, pp. 478-488; Chem. Abs., **10**, p. 2325: Experiments on the production of helium. All efforts to produce it by chemical reactions were unsuccessful.
- LO SURDO, A. Atti. Accad. Lincei (5), **30**, pp. 85-88; 1920; Chem. Abs., **15**, p. 2757: Synthetic helium and neon. Explained as contamination from atmosphere by passage through glass walls.
- MOORE, B. Nature, **90**, p. 190; 1912; Chem. Abs., **7**, p. 24: The synthesis of matter. Suggested formation of helium in X-ray bulbs.
- PIUTTI, A., and CARDOSO, E. Gazz. chim. ital., **50**, pp. 5-29; 1920; J. chim. phys., **18**, pp. 81-93; Chem. Abs., **14**, p. 2884: Experiments on the production of helium and neon in vacuum tubes containing hydrogen. Negative results.
- WYCKOFF, R. W. G. Science, **55**, pp. 130-131; 1922: A preliminary attempt to transmute lithium. Lithium ions bombarded by electrons in an attempt to produce hydrogen and helium.

VI. SEPARATION AND PURIFICATION OF HELIUM.

[For large-scale production see also Application as a balloon gas.]

- LANGLET. Z. anorg. Chem., **10**, p. 289; 1895: Preparation of helium from cleveite.
- TRAVERS. Proc. Roy. Soc., Lond., **60**, p. 449; 1896: Separation of helium from air.
- RAMSAY, W., and TRAVERS, W. Proc. Roy. Soc., Lond., **67**, pp. 329-333; 1901: Isolation of helium from the air and its physical constants.
- TRAVERS, M. W., and JAQUEROD, A. Z. physik. Chem., **45**, pp. 451-455; 1903: Note on the preparation of helium.
- DEWAR. Proc. Roy. Soc., Lond., **74**, p. 127; 1904: Separation of helium from air.
- DEWAR, J. German Patent No. 169514; 1905: Preparation of helium from air by absorption of other gases in wood charcoal at a low temperature.
- EWERS, P. Ann. Physik. (4), **17**, p. 781; 1905: Preparation and purification of helium.
- LA SOCIÉTÉ L'AIR LIQUIDE. British Patent No. 22316 of 1908: Liquefaction and rectification process for separating helium and neon from the air.
- JAQUEROD, A., and PERROT, F. J. Compt. rend., **144**, p. 135; 1907; Chem. Abs., **1**, p. 963: Preparation of pure helium by filtering the gas from cleveite through a quartz partition.
- GRAY, J. A. Proc. Roy. Soc., Lond., **82**, pp. 301-306; 1909; Chem. Abs., **3**, p. 1740: Liberation of helium from minerals by grinding.
- WOOD, D. O. Proc. Roy. Soc., Lond., **84**, pp. 70-78; 1910; Chem. Abs., **4**, p. 2082: The liberation of helium from minerals by the action of heat.

- SKAUPY, F. *Verh. deut. physik. Ges.*, **18**, pp. 230-232; 1916; *J. Chem. Sec.*, **110**, II, p. 469; 1916; *Chem. Abs.*, **10**, p. 3022: The separation of gas mixtures under the influence of the direct current. Helium and argon, among other gases, may be separated in a cathode tube.
- WEBER, S. *Ann. Physik.* (4), **54**, pp. 325-326; 437-462; 1918; *Chem. Abs.*, **13**, p. 204: Experimental researches on the conductivity of gases. Includes methods of preparation.
- EDWARDS, E., and ELWORTHY, R. T. *Trans. Roy. Soc. Can.*, **13**, III, pp. 47-52; 1919; *Chem. Abs.* **14**, p. 2531: A continuous-flow apparatus for the purification of impure helium mixtures.
- BURRELL, G. A. *Oil and Gas Journal*, **38**, pp. 50-55; 1919: Extracting helium from natural gas. Also reviews history and uses of helium.
- COTTRELL, F. G. *Chem. Met. Eng.*, **20**, pp. 104-114; 1919; *J. Ind. Eng. Chem.*, **11**, pp. 148-153; *Trans. Am. Soc. Mech. Eng.*, **41**, pp. 155-158; 1919: Commercial production of helium.
- MCLENNAN, J. C. *Trans. Roy. Soc. Can.*, **13**, III, pp. 53-56; 1919: The production of helium from the natural gases of Canada.
- SQUIER, G. O. *Proc. Am. Inst. E. E.*, **38**, p. 75; 1919; *Elec. Rev.*, **74**, p. 131; *Eng. and Min. J.*, **107**, p. 273: Production of helium on a commercial scale.
- STEWART, A. *Bur. Mines. Bull.*, **178C**, pp. 75-87; 1919: Production of helium for use in airships. In publication describing the war work of the Bureau of Mines.
- CADY, H. P., and ELSEY, H. M. K. *Science*, **50**, pp. 71-72; 1919; *Chem. Abs.*, **13**, p. 2158: The possible presence of coronium in helium from natural gas.
- DAVIS, J. W. *Eng. Mining J.*, **109**, pp. 84-85; 1920; *Chem. Abs.*, **14**, p. 2842; Methods of recovery of helium.
- BRIDEL, M. *J. pharm. chim.*, **22**, pp. 428-433; 1920: Helium, its production and uses.
- HOUSEMAN, C. R. *Chem. Age* (New York), **2**, pp. 198-199; 1920; *Chem. Trade J.*, **66**, pp. 235-236: Oxygen, nitrogen, and rare gases from the air.
- MACALLUM, A. B. Report of the administrative chairman of the Honorary Advisory Council for Scientific and Industrial Research of Canada; 1920: The conservation of helium in Canada.
- MCLENNAN, J. C. *Engineering*, **110**, pp. 154-156; 1920: Helium, its production and uses.
- MCLENNAN, J. C. *Chem. Trade J.*, **66**, pp. 827-829; 1920: Helium. War developments are described.
- MCLENNAN, J. C. *J. Chem. Soc.*, **117**, pp. 923-947; 1920: Helium, its production and uses.
- PORLEZZA, C. *Giorn. chim. ind.*, **2**, pp. 638-647, 687-691; 1920; *Chem. Abs.*, **15**, p. 2157: Industrial production of helium. A review of developments in America and of possibilities in Italy.
- SCHUETTE, O. F. *Gas Age*, **45**, pp. 192-193, 395; 1920: Production of helium.
- ELWORTHY, R. T. *Can. Chem. Met.*, **5**, pp. 7-9; 1921: History of helium and its production.
- MOORE, R. B. *J. Frank. Inst.*, **191**, pp. 145-197; 1921; *Sci. Am. M.*, **3**, p. 373: Helium, its history, properties, and commercial development.
- BAKER, B. B. *Gas Age*, **43**, p. 456; 1919: Helium, a new natural-gas by-product in demand.
- ANONYMOUS ARTICLES:
Engineering, **107**, pp. 574-575; 1919: The extraction of helium from natural gas.
Nature, **102**, pp. 487-488; 1919: The use of helium for aircraft purposes. A review of the steps taken in Great Britain for the production of helium.
Engineer, **127**, pp. 279-280; 1919: Commercial helium.
Sci. Am. Sup., **86**, p. 173; 1919: Production of helium.
Automotive Ind., **41**, pp. 578-579; 1919: History of the development of helium.
Min. and Sci. P., **120**, p. 378; 1920: Helium.

VII. ANALYSIS OF HELIUM.

- SIEVERTS, A., and BRANDT, R. *Z. angew. Chem.*, **27**, I, p. 424; 1914, and **29**, I, pp. 402-404; 1916; *Chem. Abs.*, **11**, p. 1058: Apparatus for the determination of the noble gases.
- THOMPSON, J. J. *Phil. Mag.* (6), **24**, pp. 668-672; 1912; *Chem. Abs.*, **7**, p. 17. Multiply charged atoms. Detection of helium in 1 cc of air by the positive ray method.
- ELWORTHY, R. T. *Bull.*, **31**, Mines Branch, Dept. of Mines, Canada, 1920: Describes a specific gravity balance used for determining helium.
- MURRAY, V. F. *Trans. Roy. Soc. Can.*, **13**, III, pp. 27-35; 1919: Estimation of helium in mixtures of gases by the use of a katharometer.
- McLENNAN, J. C., and ELWORTHY, R. T. *Trans. Roy. Soc. Can.*, **13**, III, pp. 19-25. 1919: Use of the Jamin interferometer for the estimation of small amounts of hydrogen or helium in the air.
- PIRANI, M., and LAX, E. *Wiss. Veröffentl. Siemens Konzern I*, pp. 167-178; 1920; *Chem. Abs.*, **15**, p. 2773: Point discharge in nitrogen and the rare gases. Development of a method for the detection of impurities in the inert gases.
- McLENNAN, J. C. *Can. Dept. Mines Bull.*, **31**, 72 pp.; 1920: Report on sources of helium in the British Empire. Contains methods of analysis.
- WEAVER, E. R., PALMER, P. E., LEDIG, P. G., PICKERING, S. F., and FRANTZ, H. W. *J. Ind. Eng. Chem.*, **12**, pp. 359-366; 1920: Automatic methods of gas analysis depending upon thermal conductivity.
- MOORE, R. B. *J. Frank. Inst.*, **191**, pp. 145-197; 1921: Helium, its history, properties, and commercial development.
- SEIBEL, C. W. *U. S. Geol. Survey. Professional Paper No. 121*; 1921: Methods of analysis of natural gas containing helium. (In a paper by G. S. Rogers entitled "Helium-bearing natural gas.")

VIII. PROPERTIES OF HELIUM.

1. ELECTRICAL AND MAGNETIC PROPERTIES.

[See also Spectrum of helium and Relation to subatomic phenomena.]

- RAMSAY and COLLIE. *Proc. Roy. Soc. Lond.*, **59**, p. 257; 1896: Behavior of argon and helium when submitted to the electrical discharge.
- STRUTT. *Phil. Mag.* (5), **49**, p. 293; 1900: Electrical discharges in argon and helium.
- RITTER. *Ann. Physik* (4), **14**, p. 118; 1904: The sparking potential in helium.
- BOUTY, E. *Compt. rend.*, **145**, p. 225; 1907; *Chem. Abs.*, **1**, p. 2853: The dielectric cohesion of helium.
- COMPTON, K. T., LILLY, E. G., and OLMSTED, P. S. *Phys. Rev.*, **16**, pp. 282-289; 1920; *Chem. Abs.*, **15**, p. 14: The minimum arcing voltage in helium.
- SODDY and MCKENZIE. *Proc. Roy. Soc. Lond.*, **80**, pp. 92-109; 1908; *Jahrb. Radioakt.*, **5**, pp. 14-17; 1908: The electric discharge in monatomic gases.
- WATSON, H. E. *Proc. Cam. Phil. Soc.*, **17**, pp. 90-107; 1912; *Chem. Abs.*, **8**, p. 289: Experiments on the electric discharge in helium and neon.
- ASTON, F. W., and WATSON, H. E. *Proc. Roy. Soc. Lond.*, **86**, pp. 168-180; 1911; *Chem. Abs.*, **6**, p. 2203: Relation between current, voltage, pressure, and the length of the dark space in different gases.
- ASTON, F. W., and KIKUCHI, T. *Proc. Roy. Soc. Lond.*, **98**, pp. 50-56; 1920; *Chem. Abs.*, **14**, p. 3592: Moving striations in neon and helium.
- MACKAY, G. M. *J. Phys. Rev.*, **15**, pp. 309-311; 1920; *Chem. Abs.*, **14**, p. 2128: Symmetrical subdivision of anode glow in helium discharge tubes.
- PERROT, A. *Compt. rend.*, **156**, pp. 1679-1681; 1912; *Chem. Abs.*, **7**, p. 3265; 1913: Movement of luminous centers in electrical discharges.

- SEELIGER, R., and POMMERENNIG, E. *Ann. Physik*, **59**, pp. 589-612; 1919; *Chem. Abs.*, **14**, p. 2130: Light emission from the cathode dark space and the negative glow boundary. Observations on nitrogen, hydrogen, oxygen, helium, and argon.
- DEFREGER, R. *Ann. Physik* (4), **12**, pp. 662-665, 1903: The cathode drop in helium.
- WATSON, H. E., and PARANJPE, G. R. *J. Indian Inst. Sci.*, **2**, pp. 43-65; 1918; *Chem. Abs.*, **14**, p. 3562: The cathode fall in neon and helium.
- KOSSEL, W. *Ann. Physik*, **37**, pp. 393-424; 1912; *Chem. Abs.*, **6**, p. 2203: Cathode rays in several gases including helium.
- FRANCK and POHL. *Verh. deut. physik. Ges.*, **9**, pp. 194-199; 1907; *Chem. Abs.*, **2**, p. 505: The mobility of the helium ion.
- NUTTING, P. G. *Bull. Bur. Stds.*, **8**, pp. 487-494; 1912; *Chem. Abs.*, **7**, p. 1140: Luminous properties of conducting helium gas.
- DORN, E. *Physik. Zeit.*, **8**, pp. 589-590; 1907; *Chem. Abs.*, **2**, p. 227: Canal rays in hydrogen, helium, and argon.
- KOENIGSBERGER and KUTSCHEWSKI. *Chem. Ztg.*, **35**, p. 616; 1911; *Chem. Abs.*, **6**, p. 2711: The behavior of helium canal rays.
- ROBINSON, J. *Physik. Zeit.*, **11**, pp. 11-13, 1910; *Chem. Abs.*, **4**, p. 863: Absorption of cathode rays by helium.
- GILL, E. W., and PIDDUCK, F. B. *Phil. Mag.* (6), **16**, pp. 280-290; 1908; *Chem. Abs.*, **3**, p. 21: Ionization by collision in argon and helium.
- GILL, E. W., and PIDDUCK, F. B. *Phil. Mag.* (6), **23**, pp. 837-849; 1912; *Chem. Abs.*, **6**, p. 2030: Ionization by collision in helium.
- BAZZONI, C. B. *Phil. Mag.* (6), **32**, pp. 566-575; 1916; *Chem. Abs.*, **11**, p. 903: Ionization potential of helium. The experimental determination does not agree with Bohr's theory.
- DAWES, H. F. *Univ. of Toronto studies. Paper No. 38 from the physical laboratories. On ionization by collision in the gases helium and argon. Toronto, the University Library*, 47 pp.; 1912.
- BEATTY, R. T. *Physik. Zeit.*, **14**, p. 622; 1913; *Chem. Abs.*, **7**, p. 3266: Ionization potentials of gases.
- COMPTON, K. T. *Phys. Rev.*, **15**, pp. 131-132; 1920; *Chem. Abs.*, **14**, p. 2582: Radiation and ionization produced in helium by 20-volt impacts.
- COMPTON, K. T. *Phil. Mag.* (6), **40**, pp. 553-569; 1920; *Chem. Abs.*, **15**, p. 465: Ionization and production of radiation by electron impacts in helium investigated by a new method.
- FRANCK, J., and KNIPPING, P. *Physik. Zeit.*, **20**, pp. 481-489; 1919: The ionizing potential of helium.
- FRANCK, J., and KNIPPING, P. *Zeit. Physik*, **1**, pp. 320-332; 1920; *Chem. Abs.*, **14**, p. 3360: Emission potentials of helium.
- FRANCK, J. *Physik. Zeit.*, **22**, pp. 388-391, 409-414, 441-448, 466-471; 1921; *Chem. Abs.*, **15**, p. 3931: The excitation of light and the ionization of atoms and molecules through the impact of slowly moving electrons.
- FOUND, C. G. *Phys. Rev.*, **15**, pp. 132-133; 1920; *Chem. Abs.*, **14**, p. 2582: Ionization potentials of argon, nitrogen, carbon monoxide, helium, hydrogen, and mercury and iodine vapors.
- FOUND, C. G. *Phys. Rev.*, **16**, pp. 41-53; 1920: Ionization potentials of several gases including helium.
- GOUCHER, F. S. *Proc. Phys. Soc. Lond.*, **33**, I, pp. 13-29; 1920; *Chem. Abs.*, **15**, p. 1920: Ionization and excitation of radiation by electron impact in helium.
- HORTON, F., and DAVIES, A. C. *Proc. Roy. Soc. Lond.*, **95**, pp. 333-353; 1919; *Chem. Abs.*, **13**, p. 1417: Investigation of the ionizing power of the positive ions from a glowing tantalum filament in helium.

- HORTON, F., and DAVIES, A. C. *Proc. Roy. Soc. Lond.*, **95**, pp. 408-429; 1919; *Chem. Abs.*, **13**, p. 1669: An experimental determination of the ionization potential for electrons in helium.
- HORTON, F., DAVIES, A. C., and BAILEY, D. *Engineering*, **108**, p. 468; 1919; *Chem. Abs.*, **14**, p. 1482: Ionization and luminosity by electron collisions.
- HORTON, F. *Phil. Mag.* (6), **39**, pp. 592-639; 1920; *Chem. Abs.*, **14**, p. 3582: Critical velocities for electrons in helium.
- HORTON, F. *Phil. Mag.* (6), **40**, pp. 440-450; 1920; *Chem. Abs.*, **15**, p. 466: The effect of a trace of impurity on the measurement of the ionization velocity for electrons in helium.
- FRANCK, J., and GROTRIAN, W. *Zeit. Physik*, **4**, pp. 89-99; 1921; *Chem. Abs.*, **15**, p. 1854: Observations on ions excited by electron impact.
- RENTSCHLER, H. C. *Phys. Rev.*, **13**, pp. 297-298; 1919; *Chem. Abs.*, **13**, pp. 3071-3072; *J. Frank. Inst.*, **188**, pp. 408-409; 1919: Resonance and ionization potentials for electrons in argon, neon, and helium.
- RENTSCHLER, H. C. *Phys. Rev.*, **14**, pp. 503-515; 1920; *Chem. Abs.*, **14**, p. 2126: Resonance and ionization potentials in argon, neon, and helium.
- STARK, J. *Jahrb. Radioakt. Elektronik*, **15**, pp. 329-364; 1918; *J. Chem. Soc.*, **116**, II, p. 262; *Chem. Abs.*, **13**, p. 3072: Ionization by canal rays.
- STEAD, G., and GOSSLING, B. S. *Phil. Mag.*, **40**, pp. 413-426; 1920; *Chem. Abs.*, **15**, p. 460: The relative ionization potentials of gases as observed in thermionic valves.
- McLENNAN, J. C., and EVANS, E. *Trans. Roy. Soc. Can.*, **14**, sec. III, pp. 19-26; 1920; *Chem. Abs.*, **15**, p. 3933: Mobilities of ions in helium at high pressure.
- MAYER, H. F. *Ann. Physik.*, **64**, pp. 451-480; 1921; *Chem. Abs.*, **16**, p. 1179: The reaction of gas molecules with free electrons of low velocity.
- RAMSAUER, C. *Ann. Physik.*, **64**, pp. 513-540; **66**, pp. 546-558; *Physik. Zeit.*, **22**, pp. 613-615; 1921; *Chem. Abs.*, **15**, p. 3933; **16**, p. 871; 1905: The effective cross-section areas of gas molecules toward slow electrons.
- HOCHHEIM, E. *Verh. deut. physik. Ges.*, **10**, pp. 446-448; 1908; *Chem. Abs.*, **2**, p. 3175: Determination of the dielectric constant of helium.
- HOCHHEIM, E. *Dissertation* (Marburg, 1909): The dielectric constant of helium.
- SKAUPY, F. *Verh. deut. physik. Ges.*, **18**, pp. 230-232; 1916; *Chem. Abs.*, **10**, p. 3022; *J. Chem. Soc.*, **110**, II, p. 469: The separation of gas mixtures under the influence of the direct current.
- SKAUPY, F. *Ber. physik. Ges.*, **19**, pp. 264-267; 1919; *Sci. Abstracts*, **22A**, p. 236; *Chem. Abs.*, **13**, p. 3072: Pressure differences in direct current discharge tubes.
- TANZLER, P. *Ann. Physik* (4), **24**, pp. 931-938; 1907; *Chem. Abs.*, **2**, p. 2180; 1908: The magnetic behavior of air, argon, and helium as compared with oxygen.

2. SPECTRUM OF HELIUM.

[See also Electrical properties and Relation to subatomic phenomena.]

- CROOKES and LOCKYER. *Compt. rend.*, **120**, p. 1103; 1895.
- CROOKES. *Chem. News*, **71**, p. 151; 1895; *Chem. News*, **72**, p. 87; 1895.
- PALMER, DE FOREST. *Am. J. Sci.* (3), **50**, p. 357; 1895.
- LOCKYER and CROOKES. *Chem. News*, **72**, p. 87; 1895.
- RAMSAY, COLLIE, and TRAVERS. *J. Chem. Soc.* **67**, p. 684; 1895.
- RUNGE and PASCHEN. *Chem. Ztg.* **19**, p. 997; 1895; *Math. naturw. Mitt. Berlin*, 1895, pp. 323, 377.
- SCHUSTER. *Chem. News*, **72**, p. 224; 1895.
- WILDE, H. *Phil. Mag.* (5), **40**, p. 466; 1895.
- TRAVERS. *Proc. Roy. Soc., Lond.*, **60**, p. 449; 1896.
- RAMSAY. *Ann. chim. phys.* (7), **13**, p. 465; 1898.

- RAMSAY and TRAVERS. *Zeit. physik. chem.* **26**, p. 135; 1898.
- LIVEING, G. D., and DEWAR, J. *Ann. chim. phys.*, **22**, pp. 482-493; 1901.
- GRAY, STEWART, HOUSTOUN, and McQUISTAN. *Proc. Roy. Soc., Lond.*, **72**, p. 16; 1903.
- LOCKYER and BAKANDALL. *Proc. Roy. Soc., Lond.*, **74**, p. 546; 1905.
- GIESEL, F. *Ber. d. chem. Ges.*; **39**, p. 2244; 1906: The spectrum of helium from radium bromide.
- SODDY, F. *Proc. Roy. Soc., Lond.*, **78**, p. 429; 1907: The spectroscopic recognition of very small amounts of helium.
- STARK, J. *Verh. deut. physik. Ges.*, **16**, pp. 468-474; 1914; *Chem. Abs.*, **8**, p. 3264: Note on the arc and spark spectrum of helium.
- MERRILL, P. W. *Bur. Stds. Sci. Paper*, No. 302; 1917; *Chem. Abs.*, **11**, p. 2559; *Bull. Bur. Stds.*, **14**, pp. 159-66; 1918: Wave lengths of the stronger lines in the helium spectrum.
- HICKS, W. M. *Trans. Roy. Soc. Lond.*, **220**, pp. 335-463; 1920; *Chem. Abs.*, **14**, p. 3590: Critical study of the spectra of the monatomic gases.
- FRANCK, J., and REICHE, F. *Zeit. Physik.*, **1**, pp. 154-160; 1920. *Chem. Abs.*, **14**, p. 2580: Helium and parhelium.
- TSCHERMACK. *Pflügers Archiv*, **88**, p. 95; 1901: Use of the helium spectrum as a standard for wave-length measurement.
- NUTTING, P. G., and TUGMAN, O. *Bull. Bur. Stds.*, **7**, pp. 49-70; 1911; *Chem. Abs.*, **6**, p. 2362. The intensities of some hydrogen, argon, and helium lines in relation to current and pressure.
- COBLENTZ, W. W. *Elec. World*, **59**, p. 365; 1912; *Chem. Abs.*, **6**, p. 1100: Special energy distribution of neon and helium.
- NICHOLSON, J. W., and MERTON, T. R. *Trans. Roy. Soc. Lond.*, **216**, pp. 459-488; 1916; *Chem. Abs.*, **10**, p. 3023: Distribution of intensity in broadened spectrum lines.
- MERTON, T. R., and NICHOLSON, J. W. *Trans. Roy. Soc., Lond.*, **226**, pp. 137-173; 1919; *Chem. Abs.*, **14**, p. 2444: Intensity relations in the spectrum of helium.
- MERTON, T. R. *Proc. Roy. Soc. Lond.*, **98**, pp. 255-260; 1920; *Chem. Abs.*, **15**, p. 631: The effect of concentration on the spectra of luminous gases.
- LYMAN, T. *Astrophysic. J.*, **33**, pp. 98-107; 1911; *Chem. Abs.*, **6**, p. 1092: The spectra of helium and several other gases in the Schumann region of short wave length.
- WOLFF, K. *Ann. Physik.*, **42**, pp. 825-839; 1913; *Chem. Abs.*, **8**, p. 859: Investigation in the extreme ultra-violet.
- RICHARDSON, O. W., and BAZZONI, C. B. *Phil. Mag.* (6), **34**, pp. 285-308; 1917; *Chem. Abs.*, **12**, p. 21: The limiting frequency in the spectra of helium, hydrogen, and mercury in the extreme ultra-violet.
- McLENNAN, J. C., and PETRIE, P. A. *Trans. Roy. Soc. Can.*, **15**, III, pp. 15-29; 1921; *Chem. Abs.*, **16**, p. 1704: The spectra of helium, hydrogen, and carbon in the extreme ultra-violet.
- FRICKE, H., and LYMAN, T. *Phil. Mag.* (6), **41**, pp. 814-817; 1921; *Chem. Abs.*, **15**, p. 2789: The spectrum of helium in the extreme ultra-violet.
- PASCHEN, F. *Ann. Physik.* (4), **27**, pp. 537-570; 1908; *Chem. Abs.*, **3**, p. 1241: Line spectra of several gases in the infra-red. Found lines for helium predicted from theory by Ritz.
- PASCHEN, F. *Ann. Physik.* (4), **36**, pp. 191-197; 1911; *Chem. Abs.*, **6**, p. 828: Accuracy of wave-length measurements in the infra-red. Redetermination of some helium lines.
- IGNATIEW, A. *Ann. Physik.* (4), **43**, pp. 1117-1136; 1914; *Chem. Abs.*, **8**, p. 3643: Interferometric wave-length measurements in the ultra-red.

- SKAUPY, F. *Zeit. Physik*, **3**, pp. 175-179; 1920; *Chem. Abs.*, **15**, pp. 799: Negative molions and the spectra of helium.
- SAHA, M. N. *Phil. Mag.* (6), **40**, pp. 72-88; 1920; *Chem. Abs.*, **15**, p. 468: Ionization in the solar chromosphere.
- SAHA, M. N. *Phil. Mag.* (6), **40**, pp. 809-824; 1920; *Chem. Abs.*, **15**, p. 799: Elements in the sun. Ionization of elements and their spectra.
- SAHA, M. N. *Phil. Mag.* (6), **41**, pp. 267-279; 1921; *Chem. Abs.*, **15**, p. 1452: Problems of temperature radiation of gases.
- RAYLEIGH. *Proc. Roy. Soc., Lond.*, **101**, pp. 114-123; 1922; *Chem. Abs.*, **16**, p. 2265: The photographic spectrum of the aurora and laboratory studies in connection with it.
- STARK, J., and GIESEL, F. *Physik. Zeit.*, **8**, pp. 580-583; 1907; *Chem. Abs.*, **2**, p. 1234: Light emission of air and helium exposed to alpha rays.
- RICHARDSON, O. W., and BAZZONI, C. B. *Nature*, **98**, p. 5; 1916; *Chem. Abs.*, **10**, p. 3024: Observations on the excitation of helium spectra.
- COMPTON, K. T., and LILLY, E. G. *Astrophys. J.*, **52**, pp. 1-7; 1920; *Chem. Abs.*, **14**, p. 3194: Excitation of the spectrum of helium. A discussion of the minimum voltage required for the excitation of the different components of the helium spectrum.
- HORTON, F., and DAVIES, A. C. *Phil. Mag.* (6), **42**, pp. 746-773; 1921; *Chem. Abs.*, **16**, p. 681: The production of radiation and ionization by electronic bombardment in pure and impure helium.
- DAVIES, A. C. *Proc. Roy. Soc., Lond.*, **100**, pp. 599-620; 1922; *Chem. Abs.*, **16**, p. 2262: Minimum electronic energies associated with the excitation of the spectra of helium.
- BIRGE, R. T. *Astrophys. J.*, **32**, 112-124; 1910; *Chem. Abs.*, **5**, p. 2593: Formulas for the spectral series for the alkali metals and helium.
- HICKS, W. M. *Trans. Roy. Soc., Lond.*, **210**, pp. 57-111; 1910; *Chem. Abs.*, **4**, p. 2903: A critical study of the spectral series; alkalies, hydrogen, and helium.
- CURTIS, W. E. *Proc. Roy. Soc., Lond.*, **89**, pp. 146-149; 1913; *Chem. Abs.*, **8**, p. 7; A new band spectrum associated with helium.
- GOLDSTEIN, E. F. *Physik. Zeit.*, **14**, pp. 624-629; 1913; *Chem. Abs.*, **7**, p. 3267: A new spectrum belonging to helium. Contains also a description of the apparatus and method of obtaining the helium used.
- KOCH, J. *Ann. Physik*, **48**, pp. 98-109; 1915; *Chem. Abs.*, **10**, p. 850: New line series in the helium spectrum.
- RIECKE. *Physik. Zeit.*, **16**, pp. 222-227; 1915; *Chem. Abs.*, **10**, 418: Bohr's theory of the series spectrum.
- FOWLER, A. *Proc. Roy. Soc., Lond.*, **91**, pp. 208-216; 1915; *Chem. Abs.*, **9**, p. 1426: New type of series in the band spectrum associated with helium.
- NICHOLSON, J. W. *Phil. Mag.* (6), **28**, pp. 90-103; 1914; *Chem. Abs.*, **8**, p. 3644: Atomic structure and the spectrum of helium.
- NICHOLSON, J. W. *Proc. Roy. Soc.*, **91**, pp. 432-439; 1915; *Chem. Abs.*, **9**, p. 2482; Band spectrum associated with helium.
- NICHOLSON, J. W. *Nature*, **94**, p. 642; 1915; *Chem. Abs.*, **9**, p. 1003: The spectra of hydrogen and helium.
- BOHR, M. *Nature*, **95**, pp. 6-7; 1915; *Chem. Abs.*, **9**, p. 1269: The spectra of hydrogen and helium.
- EVANS, E. J. *Phil. Mag.* (6), **29**, pp. 284-297; 1915: The spectra of hydrogen and helium.
- NICHOLSON, J. W. *Nature*, **95**, p. 33; 1915; *Chem. Abs.*, **9**, p. 1269: The spectra of hydrogen and helium.
- MERTON, T. R. *Nature*, **95**, p. 65; 1915; *Chem. Abs.*, **9**, p. 1269: The spectra of hydrogen and helium.

- MERTON, T. R., and NICHOLSON, J. W. Proc. Roy. Soc., Lond., **93**, pp. 27-28; 1916; Chem. Abs., **11**, p. 555: Phenomena relating to the spectra of hydrogen and helium. A study of relative intensity distribution.
- ISHIWARA, J. Proc. Tokyo Math. Phys. Soc. (2), **8**, pp. 540-550; 1916; Chem. Abs., **11**, p. 3161: Structure of the atom core. Mathematical discussion of the spectra of hydrogen and helium.
- JOHANSON, A. M. Ark. Mat. Astron. och. Fysik. Stockholm., **12**, No. 6, pp. 1-92; 1917; Sci. Abs., **22A**, p. 160; Chem. Abs., **13**, p. 3081: Series formula for the line spectra.
- LIEBERT, G. Ann. Physik, **56**, pp. 610-616; 1918; Sci. Abs., **21A**, p. 499: New helium series.
- GRENZ, W. Ber. Physik. Ges., **21**, pp. 632-643; 1919; Chem. Abs., **14**, p. 2296: Theory of band spectra in relation to molecular structure.
- KOSSEL, W., and SOMMERFELD, A. Ber. Physik. Ges., **21**, pp. 240-259; 1919; Chem. Abs., **15**, p. 799: A principle of selection and a theory of displacement in spectrum series.
- LANDE, A. Physik. Zeit., **20**, pp. 228-234; 1919; Chem. Abs., **15**, p. 18: The series spectrum of helium.
- BELL, H. Phil. Mag. (6), **40**, pp. 489-494; 1920; Chem. Abs., **15**, p. 468: The helium-hydrogen series constants.
- CURTIS, W. E. Proc. Roy. Soc. Lond., **101**, pp. 38-64; 1922; Chem. Abs., **16**, p. 2077: The structure of the band spectrum of helium.
- LYMAN, T. Nature, **104**, p. 314; 1919; Science, **50**, pp. 481-483; Chem. Abs., **14**, p. 382: A helium series in the extreme ultra-violet.
- HICKS, W. M. Nature, **104**, p. 393; 1919; Sci. Abs., **23A**, pp. 138-139; Chem. Abs., **14**, p. 3592: Helium series in the extreme ultra-violet.
- LOHMANN, W. Physik. Zeit., **9**, pp. 145-148; 1908; Chem. Abs., **2**, p. 2041: The Zeeman effect of the helium lines.
- MOLYNEAUX, JANE. Physik. Zeit., **13**, pp. 259-260; 1912; Chem. Abs., **6**, 1566: Displacement of middle component of normal Zeeman triplet in the helium spectrum.
- PURVIS, J. E. Proc. Camb. Phil. Soc., **15**, pp. 45-52; 1908; Chem. Abs., **3**, p. 1364: The rotation of various spectral lines of neon, helium, and sodium in a magnetic field.
- HANSEN, H. M., and JACOBSEN, J. C. Kgl. Danske. Videnskab. Selskab. Mathfys. Medd., **3**, No. 11, 25 pp.; 1921; Chem. Abs., **15**, p. 3941: The magnetic separation of the fine structure components of the lines of the helium spark spectrum.
- WARAN, H. P. Proc. Cambridge Phil. Soc., **20**, I, pp. 45-49; 1920; Chem. Abs., **15**, p. 338: The effect of a magnetic field on the intensity of spectrum lines: Spectra of monatomic gases greatly enhanced. Others unaffected.
- STARK. Elektrische Spektralanalyse chemischer Atome (Book), 138 pp. Leipzig, S. Hirzel; 1914: The electrical spectrum analysis of the chemical atom.
- FULCHER, G. S. Astrophys. J., **41**, pp. 359-372; 1915: The Stark effect and atomic structure.
- BRUNETTI, RITA. Atti Accad. Lincei, **24**, I, pp. 719-723; 1915; Chem. Abs., **9**, p. 2835: The Stark-Lo Surdo phenomenon in the helium spectrum.
- STARK, J. Ann. Physik, **51**, pp. 220-236; 1916; Chem. Abs., **11**, p. 232: The carrier of the principal and secondary series of the alkalis, alkaline earths, and helium. Theoretical.
- NYQUIST, H. Proc. Nat. Acad. Sci., **3**, pp. 399-401; 1917; Chem. Abs., **11**, p. 3160: The Stark effect in helium and neon.
- MERTON, T. R. Proc. Roy. Soc., Lond., **95**, pp. 30-38; 1918; Chem. Abs., **12**, p. 2492: Electric resolution and broadening of the helium lines.

- TAKAMINE, I. T., and YOSHIDA, U. *Mem. Coll. Sci. Kyoto Imp. Univ.*, **2**, pp. 325-334; 1918: The effect of an electric field on the spectrum lines of hydrogen and helium.
- TAKAMINE, I. T., and KOKUBU, N. *Proc. Tokyo Math. Phys. Soc.* (2), **9**, pp. 394-404; 1918; *Chem. Abs.*, **12**, p. 2493: Studies on the Stark effect in helium and hydrogen.
- STARK, J., HARDKE, O., and LIEBERT, G. *Ann. Physik*, **56**, pp. 569-576; 1918; *Sci. Abs.*, **21A**, p. 498; *Chem. Abs.*, **13**, p. 1558: The Stark effect for certain lines of hydrogen and helium.
- STARK, J. *Ann. Physik*, **56**, pp. 577-588; 1918; *Sci. Abs.*, **21A**, p. 498: Stark effect. New principal series of helium.
- LIEBERT, G. *Ann. Physik*, **56**, pp. 589-609; 1918; *Sci. Abs.*, **21A**, p. 499; *Chem. Abs.*, **13**, p. 1558: Stark effect for ultra-violet lines of helium.
- STARK, J. *Ann. Physik*, **60**, pp. 196-200; 1919; *Chem. Abs.*, **14**, p. 3589. The spectral evidence of electric fields in the surface of chemical atoms.
- EPSTEIN, P. S. *Ann. Physik*, **58**, pp. 553-576; 1919; *Chem. Abs.*, **14**, p. 2297: Theory of the Stark effect in Fowlers helium series.
- YOSHIDA, U. *Mem. Coll. Sci. Kyoto Imp. Univ.*, **4**, pp. 187-199; 1920; *Chem. Abs.*, **15**, p. 985: The regularity of the Stark effect on the spectral lines of hydrogen and helium.
- FOOTE, P. D. The origin of spectra. American Chem. Soc. monograph. Chemical Catalog Co., New York; 1922: Contains a critical bibliography.
- SANFORD, F. *Astrophys. J.*, **49**, pp. 337-343; 1919; *Chem. Abs.*, **13**, p. 2484: The helium spectrum and the unit electrical charge.
- GERDIEN, H., and HOLEN, R. *Ann. Physik.* (4), **27**, pp. 844-858; 1908; *Chem. Abs.*, **3**, 1242: Spectrum of the canal rays in helium.
- STARK, J. *Physik. Zeit.*, **14**, pp. 961-965; *Chem. Abs.*, **8**, p. 239; 1914: Comparison of the results of electromagnetic and spectral analysis of canal rays.
- SMITH, HAROLD. *Phil. Mag.*, **30**, I, pp. 805-811; 1915; *Chem. Abs.*, **10**, p. 422: Conductivity in mixtures of helium and hydrogen, and spectrum of the positive column.
- COLLIE. *Proc. Roy. Soc., Lond.*, **71**, p. 25; 1902: The influence of mercury vapor on the helium spectrum.
- WOOD, R. W., and FRANCK, J. *Physik. Zeit.*, **12**, pp. 81-83; 1911; *Phil. Mag.* (6), **21**, pp. 265-270; *Chem. Abs.*, **5**, 1361: Change in resonance spectrum of iodine by the presence of helium.
- WOOD, R. W. *Physik. Zeit.*, **12**, pp. 1204-1211; 1912; *Chem. Abs.*, **6**, p. 1396: Resonance spectra of iodine vapor and their destruction by gases of the helium group.
- McLENNAN, J. C., YOUNG, J. F. T., and IRETON, H. J. C. *Proc. Roy. Soc., Lond.*, **98**, pp. 95-108; 1920; *Chem. Abs.*, **15**, p. 209: Arc spectra in vacuo and spark spectra in helium of various elements.
- McLENNAN, J. C., and PETRIE, P. A. *Proc. Roy. Soc., Lond.*, **98**, pp. 109-114; 1920; *Chem. Abs.*, **15**, p. 208: Spark spectra of various elements in helium in the extreme ultra-violet.

3. OPTICAL PROPERTIES OF HELIUM.

- RAVLEIGH. *Chem. News.*, **72**, p. 223; 1895: Refractive index of helium.
- RAMSAY and TRAVERS. *Proc. Roy. Soc., Lond.*, **61**, p. 267, **62**, p. 225; 1897: Refractive index of helium. The refractive index of a mixture of equal parts of helium and hydrogen is 3 per cent lower than would be calculated from the theory of mixtures.
- RAMSAY and TRAVERS. *Proc. Roy. Soc., Lond.*, **67**, p. 329; 1900: Refractive index of helium.

- RAMSAY. Arch. Neerland Sci. Exact. Nat., **5**, p. 356; 1900: Refractivity of the inactive gases.
- SCHEEL, K., and SCHMIDT, R. Verh. deut. physik. Ges., **10**, pp. 207-210; 1908; Chem. Abs., **2**, p. 2324: The refractive index of helium.
- SCHEEL, K., and SCHMIDT, R. Physik. Zeit., **9**, pp. 921-923; 1908; Chem. Abs., **3**, p. 744: The refractive index of helium.
- HERRMANN, K. Verh. deut. physik. Ges., **10**, pp. 211-216; 1908; Chem. Abs., **2**, p. 2325: Refraction and dispersion of helium.
- HERRMANN, K. Verh. deut. physik. Ges., **10**, pp. 476-482; 1908; Chem. Abs., **2**, p. 3175: Refraction and dispersion of helium.
- BURTON, W. Proc. Roy. Soc., Lond., **80**, pp. 390-405; 1908; Chem. Abs., **3**, p. 1116: The refractive index and dispersion of light in argon and helium.
- CUTHBERTSON, C. Proc. Roy. Soc., Lond., **81**, pp. 440-448; 1908; Chem. Abs., **4**, p. 1935: The refraction and dispersion of krypton and xenon and their relation to those of helium and argon.
- CUTHBERTSON, C., and CUTHBERTSON, M. Proc. Roy. Soc., Lond., **84**, pp. 13-15; 1910; Chem. Abs., **5**, p. 232: Refraction and dispersion of argon and dispersion of helium, neon, krypton, and xenon.
- CUTHBERTSON, C., and METCALF, E. P. Proc. Roy. Soc., Lond., **80**, pp. 411-419; 1908; Chem. Abs., **3**, p. 1117: The dispersion of gaseous mercury, sulphur, phosphorus, and helium.
- DAVISSON, C. Phys. Rev., **8**, pp. 20-27; 1916; Chem. Abs., **10**, p. 2068: The dispersion of hydrogen and helium according to Bohr's theory.
- HEYDWEILER, A. Verh. deut. physik. Ges., **16**, pp. 722-727; 1914; Chem. Abs., **9**, p. 16: Atomic refraction. Includes the atomic refraction of helium in solution.
- LAYMAN, T. Astrophysic. J., **27**, pp. 87-105; 1908; Chem. Abs., **2**, p. 1657: The absorption of some gases for light of very short wave length. Helium produces no observable absorption in the region investigated.
- STRUTT, R. J. Proc. Roy. Soc., Lond., **95**, pp. 155-176; 1918: The light scattered by gases. Its polarization and intensity.
- RAYLEIGH. Nature, **104**, p. 412; 1919; Chem. Abs., **14**, p. 895: Polarization of light scattered by helium atoms.
- RAYLEIGH. Proc. Roy. Soc., Lond., **98**, pp. 57-63; 1920; Chem. Abs., **14**, p. 3591: A reexamination of the light scattered by gases in respect of polarization. Experiments on helium and argon.
- GANS, R. Ann. Physik, **65**, pp. 97-163; 1921; Chem. Abs., **15**, p. 2757: Asymmetry of gas molecules. Forms of molecules based upon the Tyndall effect.

4. SOLUBILITY AND ABSORPTION OF HELIUM.

- RAMSAY, COLLIE, and TRAVERS. J. Chem. Soc., **67**, p. 684; 1895: Solubility of helium. Entirely insoluble in alcohol and benzene. One volume of water dissolves 0.0073 volume of helium at 18.2° C.
- ESTREICHER, TH. Z. physik. Chem., **31**, p. 176; 1899: Solubility of helium in water at various temperatures.
- WINKLER, L. W. Z. physik. Chem., **55**, p. 344; 1906: Solubility of helium; questions results of Estreicher.
- ANTROPOFF, A. Proc. Roy. Soc., Lond., **83**, pp. 474-483; 1910; Chem. Abs., **4**, p. 2077: The solubility of xenon, krypton, neon, and helium in water.
- ANTROPOFF, A. Zeit. Elektrochem., **25**, pp. 269-297; 1919: Experiments on the solubility of rare gases in liquids. Helium, neon, argon, krypton, xenon, and radium emanation in water and xenon in analine. Coefficient of solubility for helium in water at 0°C-0.0097; at 50° C-0.0108.
- BERTHELOT. Compt. rend., **124**, p. 113; 1897: Supposed chemical action of helium on benzene and carbon-disulphide vapors under the electric discharge.

- DEWAR. *Comp. rend.*, **139**, p. 261; 1904: Absorption of helium in wood charcoal.
- CLAUDE, G. *Compt. rend.*, **158**, pp. 861-864; 1914; *Chem. Abs.*, **8**, p. 3734; 1914: Absorption of gases by charcoal at low temperatures. Helium, neon, hydrogen, and nitrogen investigated at liquid air temperatures.
- MCLEAN, S. *Trans. Roy. Soc. Can.*, **12**, III, pp. 79-81; 1918; *Chem. Abs.*, **13**, p. 1667: The absorption of helium by charcoal.
- TAYLOR, H. S. *J. Ind. Eng. Chem.*, **13**, pp. 75-79; 1920; *Chem. Abs.*, **15**, p. 619: Mechanism of catalytic processes. Gives results for the absorption of helium by nickel and copper.
- BRAUNER, B. *Chem. News*, **71**, p. 271; 1895: Absorption of helium in nickel electrodes.
- FRIEDLANDER, S. *Z. phys. Chem.*, **19**, p. 657; 1896: Absorption of helium in platinum electrodes of vacuum tubes.
- TRAVERS, W. M. *Proc. Roy. Soc., Lond.*, **60**, p. 449; 1896: Absorption of helium in platinum electrodes of vacuum tubes.
- STRUTT, R. J. *Proc. Roy. Soc., Lond.*, **87**, pp. 381-384; 1912; *Chem. Abs.*, **7**, p. 722: Absorption of helium and other gases under the electric discharge.
- SIEVERTS, A., and BERGNER, E. *Ber.*, **45**, pp. 2576-2583; 1912; *Chem. Abs.*, **7**, p. 2142: Solubility of argon and helium in solid and liquid metals and in quartz.
- KOHLSCHUTTER, V., and VOGT, K. *Ber.*, **38**, pp. 1419-1430, 2992-3002; 1905: Solid solutions of the inactive gases in uranium oxide.
- PIUTTI, A. *Nature*, **84**, pp. 543-544; 1910; *Chem. Abs.*, **5**, p. 626: Absorption of helium in salts and minerals.
- TROOST and OUVRARD. *Compt. rend.*, **121**, p. 394; 1895: Supposed compound of helium with magnesium.
- COOKE. *Proc. Roy. Soc., Lond.*, **77**, p. 148; 1906; *Z. phys. Chem.*, **55**, p. 537; 1906: Further researches on the chemical behavior of helium.
- RAMSAY and COLLIE. *Chem. News*, **73**, p. 259; 1896: Unsuccessful attempts to form helium compounds.

5. DENSITY AND ATOMIC WEIGHT OF HELIUM.

[See also following heading: Pressure-volume relations for gaseous helium.]

- LANGLET. *Compt. rend.*, **120**, p. 122; 1895; *Z. anorg. Chem.*, **10**, 289; 1895: Density of helium from cleveite.
- RAMSAY and TRAVERS. *Proc. Roy. Soc., Lond.*, **62**, p. 316; 1897; *Proc. Roy. Soc. Lond.*, **67**, p. 329; 1900: Good determinations of the density of helium. Ramsay's earlier determinations (not indexed) were in error.
- WATSON, H. E. *J. Chem. Soc.*, **97**, pp. 810-833; 1910; *Chem. Abs.*, **4**, p. 2076: Densities and molecular weights of neon and helium.
- International Committee on Atomic Weights, *J. A. C. S.*, **32**, 1113-1136; 1910: Atomic weight of helium.
- HEUSE, W. *Ber. physik. Ges.*, **15**, pp. 518-523; 1913; *Chem. Abs.*, **7**, p. 3864: The density and atomic weight of helium. Finds molecular weight of 4.002.
- TAYLOR, T. S. *Phys. Rev.*, **10**, pp. 653-660; 1917; *Chem. Abs.*, **12**, p. 548: Determination of the density of helium by means of a quartz microbalance.
- GUYE, P. A. *J. chim. phys.*, **16**, pp. 46-61; 1918; *Chem. Abs.*, **12**, p. 2147: Improvements to be applied to methods of weighing. The elastic microbalance. Application to the atomic weights of helium and hydrogen.
- ASTON, F. W. *Nature*, **105**, p. 8; 1920; *Chem. Abs.*, **14**, p. 1629: The constitution of the elements. Determination of the atomic weight of helium by the positive ray method.
- ASTON, F. W. *Sci. Progress*, **15**, pp. 212-222; 1920; *Chem. Abs.*, **15**, p. 335: Mass spectra and the atomic weights of the elements.
- NEWBERGER, M. C. *Physik. Zeit.*, **23**, pp. 145-146; 1922; *Chem. Abs.*, **6**, 2243: Atomic weight of helium and isohelium.

6. PRESSURE-VOLUME RELATIONS FOR GASEOUS HELIUM.

- KUENEN and RANDALL. *Chem. News*, **72**, p. 295; 1895: Pressure volume relations of helium between -210 and $+237^{\circ}$ C.
- OLSZEWSKI. *Ann. Physik. Wied.* (2), **59**, p. 184; 1896.
- RAMSAY and TRAVERS. *Proc. Roy. Soc., Lond.*, **67**, p. 329; 1900: Isotherms of helium at 11.2 and 100° C.
- TRAVERS and JAQUEROD. *Chem. News*, **86**, p. 61; 1902; *Z. Physik. Chem.*, **45**, p. 385; 1903: Coefficient of expansion of helium.
- TRAVERS and JAQUEROD. *Accad. Sci. Phys. Genève* (4), **14**, pp. 97-99; 1903: The coefficient of expansion of hydrogen and helium.
- TRAVERS and JAQUEROD. *Zeit. physik., Chem.*, **45**, pp. 456-460; 1903: The apparent value of the critical constants and boiling point of helium and an attempt to liquefy the gas.
- JAQUEROD and SCHEUER. *Compt. rend.*, **140**, p. 1384; 1905: Compressibility of helium.
- BURT, F. P. *Trans. Faraday Soc.*, **6**, pp. 19-26; 1906: The compressibility of helium and neon.
- PLANCK, I. M. *Sitz.-Ver. Akad. Wiss., Berlin*, **32**, pp. 633-647, 1909; *Chem. Abs.*, **3**, p. 132: The canonical equation of state of monatomic gases. Equation "characterizes completely the thermodynamic properties of a substance."
- GUYE, P. A., et al. *Mem. Soc. Phys. Hist. Nat., Genève.*, **35**, pp. 547-694; *Chem. Abs.*, **4**, p. 2395: Deviation of several gases from Boyles law.
- ONNES, H. K. *Proc. Acad. Amsterdam*, **10**, pp. 741-742; 1909; *Chem. Abs.*, **3**, p. 132: Isotherms of helium at -253 and -259° C.
- ONNES, H. K. *Proc. Acad. Amsterdam*, **10**, pp. 445-450; 1909; *Chem. Abs.*, **3**, p. 132: Isotherms of helium between 100 and -217° C.
- ONNES, H. K. *Verslag. k. Akad. Wet.*, **18**, pp. 168-172; *Chem. Abs.*, **5**, p. 838; 1911; Isotherms of neon and helium and their mixtures.
- ONNES, H. K. *Proc. Acad. Amsterdam*, **14**, pp. 678-684, 1913; *Chem. Abs.*, **7**, p. 2327: Isotherms of monatomic gases. Thermal properties of helium.
- ONNES, H. K. *Bull. assoc. intern. froid.*, **6**, pp. 103-134; 1916; *Chem. Abs.*, **10**, p. 1451: Brief survey of recent work of Leiden Cryogenic Laboratory containing, among other things, methods and results upon the maximum density of helium.
- HOLBORN, L., and SCHULTZE, H. *Ann. Physik*, **47**, pp. 1089-1111; 1915; *Chem. Abs.*, **10**, p. 416: Isothermals of helium. Range $0-200^{\circ}$ C. up to about 100 atmospheres.
- HOLBORN and SCHULTZE. *Z. Electrochem.*, **21**, pp. 501-511; 1915: Work of the Reichsanstalt. Pressure-volume relations of helium. Same work probably given in greater details in preceding reference.
- HENNING, F., and HEUSE, W. *Zeit. Physik.*, **5**, pp. 285-314; 1921; *Chem. Abs.*, **15**, p. 3774: Pressure and volume expansion coefficients of helium, hydrogen, and nitrogen.
- VAN LAAR, J. J. *Verslag. K. Akad. Wet.*, **24**, pp. 1635-1657; 1916; *Chem. Abs.*, **10**, p. 2817: Data on the gas constants of the helium series.
- VAN LAAR, J. J. *Zeit. anorg. allgem. Chem.*, **104**, pp. 66-76; 1918; *Chem. Abs.*, **13**, p. 3048: The additiveness of Van der Waal's constants and the fundamental values of these quantities for different substances in relation to the periodic system.
- VAN LAAR, J. J. *J. chim. phys.*, **17**, pp. 266-324; 1919; *Chem. Abs.*, **13**, p. 3045: The determination of molecular weights from density and critical constants. Atomic weight of helium is given as 4.000.
- VAN LAAR, J. J. *Chem. Weekblad*, **16**, pp. 1557-1564; 1919; *Chem. Abs.*, **14**, p. 672: The critical density of hydrogen, helium, and neon.
- VAN LAAR, J. J. *Chem. Weekblad*, **16**, pp. 1243-1250; 1919; *Chem. Abs.*, **14**, p. 138: The Avogadro normal volume and the atomic weights of hydrogen, helium, and argon.

- VAN LAAR, J. J. *Rec. trav. chim.*, **39**, pp. 215-242; 371-410; 1920: Vapor tension and heat of vaporization. Includes a discussion of the dependence of the constants of Van der Waal's equation upon temperature and volume.
- MOLLIER, A. *Physik. Zeit.*, **21**, pp. 457-463; 1920: *Chem. Abs.*, **15**, p. 1420: A graphic presentation of the equation of state.

7. DIFFUSION AND EFFUSION OF HELIUM.

- SCHMIDT, R. *Ann. Physik* (4), **14**, pp. 801-821; 1904: also separate publication: The diffusion of helium and argon.
- LONIUS, A. Dissertation (Halle, 1909): The diffusion coefficient for argon-helium mixtures and its dependence upon the composition of the mixtures.
- RAMSAY and TRAVERS. *Proc. Roy. Soc., Lond.*, **61**, p. 267; 1897: Diffusion of helium; it does not diffuse through red-hot palladium, platinum, or iron.
- JAQUEROD and PERROT. *Compt. rend.*, **139**, p. 789; 1904: The diffusion of helium through hot quartz. Its use as a thermometric substance.
- JAQUEROD and PERROT. *Arch. sci. phys. nat.* (4), **18**, pp. 613-615; 1904: *Arch. sci. phys. nat.* (4), **20**, p. 128; 1905: The diffusion of helium through silica at high temperatures.
- DEWAR, J. *Proc. Roy. Inst.*, **21**, pp. 813-826; 1918: *J. Chem. Soc.*, **114**, II, pp. 186-187; *Chem. Abs.*, **12**, p. 2275: Diffusion of gases through rubber.
- McLENNAN, J. C., and SHAVER, W. W. *Phil. Mag.* (6), **40**, pp. 672-681; 1920; *Chem. Abs.*, **14**, p. 3568: The permeability of thin fabrics to hydrogen and helium.
- ELWORTHY, R. T., and MURRAY, V. F. *Trans. Roy. Soc. Can.*, **13**, III, pp. 37-45; 1919; *Chem. Abs.*, **14**, p. 2555: Permeability of balloon fabrics to hydrogen and to helium.
- EDWARDS, J. D., and PICKERING, S. F. *Bur. Stds. Sci. Paper*, No. 387; 1920; *Chem. Met. Eng.*, **23**, pp. 17-21; 71-75; *Chem. Abs.*, **14**, p. 2733: Permeability of rubber to gases.
- RAMSAY. *Ann. chim. phys.* (7), **13**, p. 433; 1898: Effusion of helium. It is 10 per cent greater relative to hydrogen than the difference in densities would indicate.
- DONNAN. *Phil. Mag.* (5), **49**, p. 423; 1900: Viscosity of helium; rate of effusion.

8. VISCOSITY OF HELIUM.

[See also preceding heading: Effusion of helium.]

- RAYLEIGH. *Proc. Roy. Soc., Lond.*, **59**, p. 198; 1895: Viscosity of helium.
- SCHULTZE, H. *Ann. Physik* (4), **6**, pp. 302-314; 1901: The internal friction of helium and its alteration with temperature.
- TANZLER, P. *Ber. physik. Ges.* (4), p. 222; 1906: The coefficient of internal friction of helium.
- SCHIERLOH. Dissertation (Halle; 1908): The coefficient of internal friction of argon and helium.
- RANKINE, A. O. *Proc. Roy. Soc., Lond.*, **83**, pp. 265-276; 516-535; 1910; *Physik. Zeit.*, **11**, pp. 497-502; *Chem. Abs.*, **4**, pp. 2223, 3155: Viscosity of the gases of the argon group.
- RANKINE, A. O. *Proc. Roy. Soc., Lond.*, **84**, pp. 181-192; 1910; *Physik. Zeit.*, **11**, pp. 745-752; *Chem. Abs.*, **4**, p. 3155: The variation with temperature of the viscosities of the gases of the argon group.
- ONNES, DORSMAN, and WEBER. *Verslag. K. Akad. Wet.*, **21**, pp. 1375-1384, 1385-1391, 1530-1534; 1913; *Proc. Acad. Amsterdam*, **15**, pp. 1386-1403; 1913; *Leiden Comm. Physic. Lab.*, **134**, a, b, c: Investigation of the viscosity of gases at low temperatures.
- GILÉ, A. *Ann. Physik.*, **48**, pp. 799-837; 1915; *Chem. Abs.*, **10**, p. 844: Viscosity coefficients of mixtures of hydrogen and helium.

NERNST, W. Sitz.-Ber. preuss. Akad., 1919, pp. 118-127; Chem. Abs., 13, p. 1962: Deductions from the degeneration theory of gases. Includes a derivation of the internal friction of helium.

HOFSSASS, M. J. Gasbel., 62, pp. 776-777; 1919: Viscosity of gases.

9. SPECIFIC HEATS AND OTHER THERMAL PROPERTIES OF HELIUM.

LANGLET. Zeit. anorg. Chem., 10, p. 289; 1895: Ratio of specific heats of helium.

RAMSAY, COLLIE, and TRAVERS. J. Chem. Soc., 67, p. 684; 1895: Ratio of specific heats of helium.

THOMAS. Dissertation (Marburg; 1905): Determination of the specific heat of helium. The atomic heat of helium and argon.

BEHN, U., and GEIGER, H. Verh. deut. physik. Ges., 9, pp. 657-666; 1907; Chem. Abs., 2, 1220: A modification of the Kundt method of determining the specific heat ratio and a new determination of this ratio for helium.

EGGERT. Dissertation (Marburg; 1910): Determination of specific heat of helium at various temperatures and its significance in connection with the kinetic theory of monatomic gases.

SCHELL, K., and HEUSE, W. Sitz.-Ber. Akad., Berlin, 1913, pp. 44-48; Chem. Abs., 7, p. 2507. Specific heat of helium and of certain diatomic gases between 20 and 180°.

EGGERT, A. Ann. Physik., 44, pp. 643-656; 1914; Chem. Abs., 8, p. 2977: Determination of the specific heat of helium at ordinary and higher temperatures. No indication that specific heat changes with temperature.

EUCKEN, A. Verh. deut. physik. Ges., 18, pp. 4-17; 1916; Chem. Abs., 10, p. 1129: Thermal behavior of some compressed and condensed gases at low temperatures. Specific heats of hydrogen (gas at several pressures and liquid), helium, argon (solid and liquid), nitrogen (liquid and two solid forms), oxygen (liquid and three solid forms), carbon monoxide (liquid and two solid forms), and carbon dioxide (solid at various temperatures). Heat of fusion, transformation, and evaporation of the various phases.

ONNES, H. K. Proc. Acad. Amsterdam., 14, pp. 678-684; 1911; Chem. Abs., 7, p. 2327: Thermal properties of helium.

EUCKEN, A. Sitz.-Ber. preuss. Akad. 1914, pp. 682-693; Chem. Abs., 8, p. 3141: Quantum effect with monatomic gases and liquids. Atomic heat of compressed helium at very low temperature.

HARBUTT, J. Phys. Zeit., 22, pp. 52-53; 1921; Chem. Abs., 15, p. 1237; Sci. Abs., 24, p. 431: An approximate calculation of the heat of fusion of the inert gases. Value for helium less than for any other known substance.

10. THERMAL CONDUCTIVITY OF HELIUM.

SCHWARZE, W. Physik. Zeit., 4, p. 229; 1903: The heat conductivity of argon and helium.

SCHWARZE, W. Ann. Physik. (4), 11, pp. 303-330; 1903: Determination of the thermal conductivity of argon and helium by the method of Schiermacher.

EUCKEN, A. Physik. Zeit., 13, pp. 1101-1107; 1912; Chem. Abs., 6, p. 1087; 1912: The dependence on temperature of the heat conductivity of certain gases.

WEBER, S. Ann. Physik., 54, pp. 325-326; 437-462; 1918; Sci. Abs., 21A, pp. 268-269; Chem. Abs., 13, p. 204: Experimental researches on the conductivity of gases. Includes methods of preparing helium, etc.

WEBER, S. Ann. Physik, 54, pp. 481-502; 1918; Sci. Abs., 21A, pp. 268-269; Chem. Abs., 13, p. 204: Theoretical and experimental researches on the thermal conductivity of gas mixtures.

IX. LIQUEFACTION AND PROPERTIES OF LIQUID HELIUM.

- OLSZEWSKI, K. *Ann. Phys. Wied.* (2), **59**, p. 184; 1896: Attempt to liquefy helium.
- DEWAR. *Proc. Chem. Soc., Lond.*, pp. 129 and 195, 1897-98: Attempt to liquefy helium.
- DEWAR. *Chem. News*, **84**, p. 49; 1901: Attempt to liquefy helium.
- DEWAR. *Proc. Roy. Soc., Lond.*, **68**, pp. 360-366; 1901: Experiments on the liquefaction of helium at the temperature of liquid hydrogen.
- DEWAR. *Compt. rend.*, **139**, pp. 421-422; 1904: Liquefaction of helium.
- TRAVERS and JAQUEROD. *Z. phys. Chem.*, **45**, pp. 456-460; 1903: Attempt to liquefy helium.
- OLSZEWSKI. *Z. komprim. Gase*, **9**, pp. 54-56; 1905; *Ann. Physik.* (4), **17**, p. 944, 1905; *Ann. chim. phys.* (8), **8**, pp. 139-144; 1906: Attempt to liquefy helium. Author believed he had reached a temperature of 1.7° absolute without success.
- DEWAR, J. *Engineering*, **85**, p. 793; 1908: Nadir of temperature and allied problems. Abstract of address at the Royal Institution. Discusses the difficulties attending the liquefaction of helium.
- ONNES, H. K. *Verslag. k. Akad. Wet.*, **16**, pp. 819-822; 1908; *Proc. Acad. Amsterdam*, **10**, p. 744; 1908. *Chem. Abs.*, **3**, 132: Experiments on the condensation of helium by expansion.
- ONNES, H. K. *Chem. Ztg.*, **32**, pp. 901-903; 1908; *Chem. Abs.*, **2**, 3317: The liquefaction of helium. Contains also some physical constants of the element.
- ONNES, H. K. *Verslag. k. Akad. Wet.*, **17**, pp. 163-179; 1908; *Proc. Acad. Amsterdam*, **11**, pp. 168-185, 1908; *Leiden Comm. Phys. Lab.*, No. 108, 1908; *Arch. néer. Sci. Soc.* (2), **14**, pp. 289-308; *Chem. Abs.*, **4**, p. 550: The liquefaction of helium.
- ONNES, H. K. *Compt. rend.*, **147**, pp. 421-424; 1908: Liquid helium.
- ANONYMOUS. *Science*, **28**, pp. 180, 316; 1908: Liquefaction of helium; reference to work of Onnes.
- ANONYMOUS. *Chem. News*, **98**, p. 37; 1908: Liquefaction of helium; reference to work of Onnes.
- LORENTZ, H. A. *Arch. néerland. Sci.* (2), **13**, p. 492, 1909; *Chem. Abs.*, **3**, p. 1848: Liquefaction of helium. The history and achievements of the Onnes laboratory in Leyden to the liquefaction of helium.
- ONNES, H. K. *Chem. Ztg.*, **34**, pp. 1373-1374; 1910; *Chem. Abs.*, **5**, p. 1010: Investigations carried out in the low-temperature laboratory at Leiden.
- ONNES, H. K. *Proc. Acad. Amsterdam*, **13**, pp. 1093-1113; 1911; *Chem. Abs.*, **6**, p. 6: Experiments with liquid helium. Liquid-helium bath, constant-volume helium thermometer, vapor density, vapor pressure, and density of liquid helium.
- ONNES, H. K. *Proc. Acad. Amsterdam*, **14**, pp. 204-210; 1911; *Chem. Abs.*, **7**, p. 2327: Experiments with liquid helium. A helium cryostat.
- ONNES, H. K., and WEBER, S. *Proc. Acad. Amsterdam*, **18**, pp. 493-507; 1915; *Chem. Abs.*, **10**, p. 308: Vapor pressure of helium. Temperature obtained with liquid helium. Discussion of low-temperature measurement.
- MEISSNER. *Zeit. Electrochem.*, **21**, pp. 501-511; 1915: Work of Reichsanstalt on the liquefaction of hydrogen and helium.
- MEISSNER, W. *Ber. physik. Ges.*, **21**, pp. 369-382; 1919; *Chem. Abs.*, **14**, p. 680; *Sci. Abs.*, **22A**, p. 482: Electric work during fluid emission. An improved method for the liquefaction of gases.

X. APPLICATIONS OF HELIUM.

1. APPLICATION OF HELIUM TO THERMOMETRY.

- KUENEN and RANDALL. *Chem. News*, **72**, p. 295; 1895: Use of helium in low-temperature thermometers.
- OLZEWSKI. *Ann. Phys. Wied.* (2), **59**, p. 184; 1896: Use of helium in low-temperature thermometers.
- DEWAR. *Am. J. Sci.* (4), **11**, p. 291; 1901: Use of helium for low-temperature thermometers.
- TRAVERS, SENTER, and JAQUEROD. *Chem. News*, **86**, p. 61; 1902: Use of helium in low-temperature thermometers.
- ONNES, H. K. *Proc. Acad. Amsterdam*, **10**, pp. 589-591; 1908; *Chem. Abs.*, **3**, p. 132: Derivation of the pressure coefficient of helium for the international thermometer and the reduction of the readings of the helium thermometer to the absolute scale.
- ONNES, H. K., and HOLST, G. *Verslag. k. Akad. Wet.*, **23**, pp. 175-182; 1914; *Chem. Abs.*, **9**, p. 747: The measurement of very low temperatures. A comparison of the hydrogen, helium, and platinum resistance thermometers at temperatures down to the freezing point of hydrogen.
- HOLBORN, L., and HENNING, F. *Ann. Physik.*, **35**, pp. 761-764; 1911; *Chem. Abs.*, **5**, p. 3642: Comparison of the platinum, nitrogen, hydrogen, and helium thermometers and the determination of certain fixed points between 200 and 450°.
- CATH, P. G., ONNES, H. K., and BURGERS, J. M. *Proc. Acad. Sci. Amsterdam*, **20**, pp. 1163-1176; 1918; *Chem. Abs.*, **12**, p. 2267: Comparison of the platinum and gold resistance thermometers with the helium thermometer.
- JAQUEROD and PERROT. *Compt. rend.*, **139**, p. 789, 1904: Diffusion of helium through quartz at high temperatures makes it inapplicable for use in thermometers for measuring high temperatures.
- DORN, E. *Physik. Zeit.*, **7**, p. 312; 1906: Use of helium in a platinum-iridium vessel to measure high temperatures.
- CARIO. *Dissertation* (Halle, 1907): Use of helium in platinum-iridium vessels at high temperatures.

2. APPLICATIONS OF LIQUID HELIUM ESPECIALLY TO DETERMINATION OF ELECTRICAL PROPERTIES OF METALS AT LOW TEMPERATURES.

[See also Liquefaction and properties of liquid helium.]

- ONNES, H. K., is the author of all articles under this heading unless otherwise noted.
- Leiden, Van Bemmelen *Gedenkboek*, pp. 441-446; *Chem. Abs.*, **5**, p. 3186: The attainment of temperatures considerably below the boiling point of helium.
- Electrician*, **67**, p. 657, 1911; *Chem. Abs.*, **5**, p. 3541: Electrical resistance of pure metals at liquid helium temperatures.
- Proc. Acad. Amsterdam*, **13**, pp. 1274-1276; 1911; *Chem. Abs.*, **6**, p. 6: Resistance of pure mercury at liquid helium temperatures.
- Proc. Acad. Amsterdam*, **14**, pp. 113-115; 1911; *Chem. Abs.*, **6**, p. 6; **7**, p. 2327: The electrical resistance of pure metals. The sudden change in the rate at which the resistance of mercury disappears.
- Electrician*, **71**, pp. 855-857, *Chem. Abs.*, **7**, p. 3924; 1913: Electrical resistance of pure metals at liquid helium temperatures.
- Verslag. k. Akad. Wet.*, **1913**, pp. 1284-1305, 1388-1391; *Chem. Abs.*, **8**, p. 1907; 1914: The potential difference necessary for the flow of an electric current through mercury below 4.19° absolute.
- Verslag. k. Akad. Wet.*, **22**, pp. 1413-1421; **23**, pp. 167-172; 1914; *Chem. Abs.*, **9**, p. 545: Imitating a permanent magnet by means of superconductors.

- Comp. rend., **159**, pp. 34-38; 1914; Chem. Abs., **8**, p. 3264: The persistence of electric currents without electromotive force in the superconductors.
- Proc. Acad. Amsterdam, **16**, pp. 673-688; 1914; Chem. Abs., **8**, p. 2294: The sudden disappearance of the ordinary resistance of tin and the superconductive state of lead.
- Proc. Acad. Amsterdam, **16**, pp. 987-992; 1914; Chem. Abs., **9**, p. 735: Hall effect and the magnetic change in resistance at low temperature. The appearance of galvanic resistance in supraconductors which are brought into a magnetic field at the threshold value of the field.
- Verslag. k. Akad. Wet., **22**, pp. 1027-1033; Chem. Abs., **8**, p. 2107; 1914: Hall effect and the change of resistance in a magnetic field at low temperatures.
- Verslag. k. Akad. Wet., **23**, pp. 172-175; 1914; Chem. Abs., **9**, p. 545: Phenomena of incipient paramagnetic saturation.
- Verslag. k. Akad. Wet., **23**, pp. 493-501; 1914: Hall effect and the change of resistance in magnetic fields.
- Verslag. k. Akad. Wet., **23**, pp. 703-710; 1914; Chem. Abs., **9**, p. 1580: Specific heat and thermal conductivity of mercury and some measurements of the conductivity and thermoelectromotive force at liquid-helium temperatures.
- ONNES, DORSMAN, and WEBER. Verslag. k. Akad. Wet., **21**, pp. 1375-1384, 1385-1391, 1530-1534; 1913; Proc. Acad. Amsterdam **15**, pp. 1386-1403; 1913; Leiden Comm. Phys. Lab., **134**, a, b, c.: Investigation of the viscosity of gases at low temperatures.
- Bull. assoc. intern. froid., **6**, pp. 103-134; 1915; Chem. Abs., **10**, p. 1451: Report on researches made in the Leiden Cryogenic Laboratory between the second and third congresses of refrigeration. A review.

3. APPLICATIONS OF HELIUM TO PHOTOMETRY, ETC.

- TSCHERMAK, A. Pflügers, Archiv, **88**, p. 95; 1901: Use of the helium spectrum as a standard for wave-length measurement.
- NUTTING, P. G. Bull. Bur. Stds., **4**, pp. 511-523; 1908; Chem. Abs., **2**, p. 2895: The helium tube as a primary light standard.
- NUTTING, P. G. J. Wash. Acad. Sci., **1**, pp. 221-223; 1911; Chem. Abs., **6**, p. 543; Elec. Rev. West Elec., **59**, p. 1074: Helium tubes as light standards.
- HIRSHBERG, L. K. J. Gas. Ltg., **118**, p. 720; 1912; Chem. Abs., **6**, p. 3002: Helium as a unit of candlepower.
- DORN, E. Ann. Physik, (4), **16**, p. 784; 1904: Suggested application of helium-filled Plücker tube for the recognition of rapid electrical fluctuations.
- ZEHNDER, L. Physik. Zeit., **13**, pp. 446-449; 1912: Use of vacuum tubes containing helium to demonstrate properties of electric waves.

4. APPLICATION OF HELIUM AS A BALLOON GAS.

- BURRELL, G. A. Oil and Gas Journal, **38**, pp. 50, 55; 1919: Extracting helium from natural gas. Also reviews history and uses of helium.
- COTTRELL, F. G. Chem. Met. Eng., **20**, pp. 104-114; 1919; J. Ind. Eng. Chem., **11**, pp. 148-153; Am. Soc. Mech. Eng., **41**, pp. 155-158: Commercial production of helium. Gives history of its development as a balloon gas.
- ROGERS, G. S. Nat. Geog. Mag., **35**, pp. 441-456; 1919: Helium, the new balloon gas.
- MOORE, R. B. Proc. Am. Mining Congress, **22**, p. 781; 1919: Development of helium.
- SQUIER, G. O. Proc. Am. Inst. E. E., **38**, p. 75; 1919; Trans. A. I. E. E., **38**, p. 23: Production of helium on a commercial scale.
- STEWART, A. Bur. Mines Bull., **178C**, pp. 75-87; 1919: Production of helium for use in airships.

- STEWART, A. *Aviation*, **7**, pp. 32-35; 1919: Production of helium for use in airships. U. S. AIR SERVICE (War Department): *Airship and balloon gas manual* (1919).
- ANON. *Nature*, **102**, pp. 487-488; 1919: The use of helium for aircraft purposes. A review of the steps taken in Great Britain for the production of helium.
- BLANCHARD, H. F. *Automotive Ind.*, **40**, pp. 845-849; 1919: Inauguration of safe aerial transportation.
- D'ORCY, L. *Aviation*, **5**, pp. 695-697; 1919: Future of the helium airship.
- D'ORCY, L. *Sci. Am.*, **120**, p. 73; 1919: Final solution of the airship problem.
- ANON. *Engineer*, **127**, pp. 279-280: Commercial helium.
- ANON. *Gas Age*, **44**, p. 155; 1919: Helium.
- BRIDEL, M. *J. pharm. chim.*, **22**, pp. 428-433; 1920: Helium, its production and uses.
- ANDREWS, W. S. *Gen. Elec. Rev.*, **23**, pp. 227-228; 1920: Helium, the substitute for hydrogen in airships.
- McLENNAN, J. C. *J. Chem. Soc.*, **117**, pp. 923-947; 1920: Engineering, **110**, pp. 154-156; 1920: Helium, its production and uses.
- ANON. *Sci. Am.*, **123**, p. 493; 1920: Uses of helium.
- MOORE, R. B. *J. Frank. Inst.*, **191**, pp. 145-197; 1921: Helium, its history, properties, and commercial development.
- SKERRETT, R. G. *Compressed Air Magazine*, **23**, pp. 10272-10276; 1921: Why America leads in production of helium.
- LITTLE, H. P. *Sci. Am. Monthly*, **4**, pp. 177-178; 1921: Balloons, helium gas, and the age of the earth.
- ANON. *Aerial Age*, **14**, p. 203; 1921: Helium for airships.
- ANON. *Brass World and Platers Guide*, **17**, pp. 327-328; 1921: Helium.
- ANON. *Aviation*, **11**, pp. 635-636; 1921: Helium for airships.
- ANON. *Aerial Age*, **14**, p. 316; 1921: Helium a success in airship tests.
- MOORE, R. B. *Bur. of Mines, Serial No. 2363*; 1922: Helium. Report of investigations of Bureau of Mines.
- MOORE, R. B. *Natural Gas*, **111**, p. 14, 1922: Helium, its history, properties, and commercial development.
- MOORE, R. B. *The Carolina Chemist*, **8**, p. 7; 1922: Helium.
- ANON. *Sci. Am. M.*, **1**, p. 330; 1920: Industrial applications of helium.
- SATTERLY, J., and BURTON, E. F. *Trans. Roy. Soc. Can.*, **13**, III, pp. 211-215; 1919: The combustibility of mixtures of hydrogen and helium.
- LEDIG, P. G. *J. Ind. Eng. Chem.*, **12**, pp. 1098-1103; 1920: Inflammability of jets of hydrogen and inert gas. (Nitrogen and helium.)

XI. RELATION OF HELIUM TO SUBATOMIC PHENOMENA.

[See also Formation of helium, Electrical and magnetic properties, and Spectrum of helium.]

- ERFLE, H. *Ver. deut. physik. Ges.*, **10**, pp. 331-338; 1908; *Chem. Abs.*, **2**, p. 2493: The number of quasi-bound electrons in the helium atom.
- CREHORE, A. C. *Phys. Rev.*, **34**, pp. 247-258; 1912; *Chem. Abs.*, **6**, p. 2206: Theory of atomic structure and calculation of some atomic weights.
- LORING, F. H. *Chem. News*, **106**, p. 37; 1912; *Chem. Abs.*, **6**, p. 3050: Is helium fundamentally an element of electropositive make-up?
- PERRIN, J. P. *Atoms*. French edition, 1913. English edition, 1916; New York; D. Van Nostrand Co.
- TAYLOR, T. S. *Phil. Mag.* (6), **26**, pp. 402-410; 1913; *Chem. Abs.*, **7**, p. 3710; 1913: The range and ionization of the alpha particles in simple gases.
- WILDE, H. *Chem. News*, **108**, p. 25; 1913; *Chem. Abs.*, **7**, p. 3432: Some new relations of atomic weights and transformation of neon and helium.

- HARKINS, W. D., and WILSON, E. D. *J. Am. Chem. Soc.*, **37**, pp. 1383-1396; 1915; *Chem. Abs.*, **9**, p. 2335: Structure of complex atoms; the hydrogen-helium system. Accounts for composition of 26 elements.
- PARSON, A. L. A magnetic theory of the structure of the atom. 80 pp. Smithsonian Institution; 1915.
- RUTHERFORD, E. *Smithsonian Inst. Ann. Rep.*, 1915, pp. 167-202: The constitution, of matter and the evolution of the elements.
- CREHORE, A. C. *The mystery of matter and energy*. New York, D. Van Nostrand Co., 1917.
- CREHORE, A. C. *Proc. Am. Inst. Elec. Eng.*, **37**, pp. 283-315; 1918; *Chem. Abs.*, **13**, p. 1414: Some applications of electromagnetic theory to matter.
- KUNZ, J. *Phys. Rev.*, **12**, pp. 59-69; 1918; *Chem. Abs.*, **12**, p. 2160: Bohr's atom and magnetism. Theoretical, based on behavior of hydrogen and helium.
- SCHMIDT, C. *Zeit. anorg. allgem. Chem.*, **103**, pp. 79-118; 1918; *Chem. Abs.*, **13**, p. 2623: Periodic system and generation of the elements.
- SOMMERFIELD, A. *Ann. Physik*, **53**, pp. 497-550; 1918; *Chem. Abs.*, **12**, p. 2159: The probable constitution of the helium atom (included in a theoretical paper of different title).
- WENZEL, F. *J. prakt. Chem.*, **98**, pp. 155-203; 1918; *Chem. Abs.*, **14**, p. 2292: The chemical structure of atoms.
- GEHRCKE, E. *Ber. physik. Ges.*, **21**, pp. 779-784; 1919: The structure of atomic nuclei.
- HACKH, I. W. D. *Science*, **49**, pp. 328-329; 1919; *Chem. Abs.*, **13**, p. 1414: Old age of chemical elements. Predicts the occurrence of isotopes of helium.
- LANGMUIR, I. *Proc. Am. Acad. Sci.*, **5**, pp. 252-259; 1919: The structure of atoms and the octet theory of valence.
- LANGMUIR, I. *J. Am. Chem. Soc.*, **41**, pp. 868-934; 1919; *Chem. Abs.*, **13**, p. 1554: The arrangement of electrons in atoms and molecules.
- THOMSON, J. J. *Engineering*, **107**, pp. 366-367, 410-411, 443-444, 476-478, 511-512, 560-562; 1919; *Electrician*, **82**, pp. 380, 473, 497, 519, 535: Spectrum analysis and atomic structure. Report of a series of six lectures before the Royal Institution.
- THOMSON, J. J. *Rev. sci.*, **57**, pp. 129-130; 1919; *Chem. Abs.*, **14**, p. 2292: The transmutation of the elements.
- CREHORE, A. C. *J. Elec.*, **44**, pp. 325-326; 1920: Planetary attraction and the helium atom.
- DUSHMAN, S. *J. Am. Inst. Elec. Eng.*, **39**, pp. 119-131; 1920: A review of present theories of atomic structure.
- FEHRLE, K. *Physik. Zeit.*, **21**, 552-554; 1920; *Chem. Abs.*, **15**, p. 195: A resonance theory of chemical combination.
- HARKINS, W. D. *Nature*, **105**, pp. 230-231; 1920; *Chem. Abs.*, **14**, p. 2123: Hypothetical discussion of the constitution of the atom.
- HINSBERG, O. *J. prakt. Chem.*, **101**, pp. 97-112; 1920; *Chem. Abs.*, **15**, p. 624: Binuclear theory of the atom and the periodic system.
- KEMBLE, E. C. *Science*, **52**, pp. 581-583; 1920; *Chem. Abs.*, **15**, p. 182: Helium and hydrogen molecules.
- KOHLWEILER, E. *Zeit. physik. Chem.*, **94**, 513-541; 1920; *Chem. Abs.*, **14**, p. 2747: Constitution and configuration of the atom.
- LADENBURG, R. *Zeit. Elektrochem.*, **26**, pp. 262-274; 1920; *Chem. Abs.*, **14**, p. 3187: Atomic structure and the periodic system of the elements.
- LANDE, A. *Physik. Zeit.*, **21**, pp. 114-120; 1920; *Chem. Abs.*, **15**, p. 19: Disturbance theory of the helium atom.

- LANGMUIR, I. *Science*, **51**, pp. 605-607; 1920; *Chem. Abs.*, **14**, p. 2580: The structure of the helium atom.
- LENZ, W. *Naturwissenschaften*, **8**, pp. 181-186; 1920; *Chem. Abs.*, **14**, p. 3571: Rutherford's experiments on the subdivision of the nitrogen nucleus.
- LORING, F. H. *Chem. News*, **120**, pp. 181-184, 193-196, 205-207, 217-219, **121**, pp. 2-3; 1920; *Chem. Abs.*, **14**, p. 2580: The structure of matter and the quantum theory.
- LORING, F. *Atomic Theories*; 218 pp. 1921. London, Methuen & Co.
- RUTHERFORD, E. E. *Proc. Roy. Soc. Lond.*, **97**, pp. 374-401; 1920; *Chem. Abs.*, **14**, 3570: Nuclear constitution of atoms.
- SOMMERFELD, A. *Zeit. Elektrochem.*, **26**, pp. 258-260; 1920; *Chem. Abs.*, **14**, p. 3187: The foundation and aim of the Bohr theory of atoms and spectra.
- STARK, J. *Jahrb. Radioakt. Elektronik*, **17**, pp. 161-173; 1920: Criticism of Bohr's theory of the emission of light.
- BOTHE, W. *Zeit. Physik*, **4**, pp. 300-314; 1921; *Chem. Abs.*, **15**, p. 2240. Theory of the scattering of alpha particles at small angles.
- BROSSLERA, F. *Rev. chim.*, **1**, pp. 42-48, 74-80; 1921; *Chem. Abs.*, **15**, p. 358: The existence of helium nuclei in the nuclei of radioactive elements.
- CHWOLSON, O. *Zeit. physik*, **7**, pp. 268-284; 1921; *Chem. Abs.*, **16**, p. 2067. The structure of the atomic nucleus.
- DINGLE, H. *Sci. Progress*, **16**, pp. 223-224; 1921: Significance of spectroscopy. The various types of spectra and their bearing on the ultimate structure of matter.
- HARKINS, W. D. *Phil. Mag.* (6), **42**, pp. 305-339; 1921; *Chem. Abs.*, **16**, p. 16: Constitution and stability of atom nuclei.
- KEMBLE, E. C. *Phil. Mag.* (6), **42**, pp. 123-133; 1921; *Chem. Abs.*, **15**, p. 2775: The probable normal state of the helium atom.
- LANGMUIR, I. *Phys. Rev.*, **17**, pp. 339-353; 1921; *Chem. Abs.*, **15**, p. 238: The structure of the helium atom.
- LORENTZ, H. A. *Physica*, **1**, pp. 228-241; 1921: The theoretical importance of the Zeeman effect. A review of the theories developed to relate the Zeeman effect to the structure of the atom.
- MILLIKAN, R. A. *Phys. Rev.*, **18**, pp. 456-460; 1921; *Chem. Abs.*, **16**, p. 1041: Some facts bearing on the structure of atoms, particularly the helium atom.
- NEUBURGER, M. C. *Zeit. physik. Chem.*, **99**, pp. 454-473; 1921: A new nuclear model.
- PEASE, R. N. *J. Am. Chem. Soc.*, **42**, pp. 991-1004; 1921; *Chem. Abs.*, **15**, p. 2570: An analysis of molecular volumes from the point of view of the Lewis-Langmuir theory of molecular structure.
- RUTHERFORD, E., and CHADWICK, J. *Phil. Mag.*, **42**, pp. 809-825; 1921; *Chem. Abs.*, **16**, p. 526: Artificial disintegration of light elements.
- SMEKAL, A. *Naturwissenschaften*, **9**, pp. 77-82, 93-98; 1921: Rutherford's discovery of a new light-atom nucleus.
- WEBSTER, D. L., and PAGE, L. *Bull. Natl. Research Council*, **2**, pp. 336-395; 1921: General survey of the present status of the atomic structure problem.
- BOHR, N. *Zeit. Physik*, **9**, pp. 1-67; 1922; *Chem. Abs.*, **16**, p. 2252: Atomic structure and the physical and chemical properties of the elements.
- CREHORE, A. C. *Phil. Mag.* (6), **43**, pp. 886-914; 1922; *Chem. Abs.*, **16**, p. 2448: Atoms and molecules.
- HARKINS, W. D., and MADORSKY, S. L. *Phys. Rev.*, **19**, pp. 135-156; 1922; *Chem. Abs.*, **16**, p. 1358: A graphical study of the stability relations of atom nuclei.
- MCLENNAN, J. C. *Science*, **55**, pp. 219-232; 1922: Atomic nuclei. An address discussing isotopes, and the size, electric charge, fields of force, and structure of atomic nuclei.

XII. MISCELLANEOUS AND GENERAL.

- RAMSAY, WILLIAM. *Proc. Roy. Soc.*, **60**, p. 53; 1896: Helium and argon. Experiments which show the inactivity of these elements.
- RAMSAY, WILLIAM. *Ann. Rep. Smithsonian Inst.* 1898, pp. 267-276: The recently discovered gases and their relation to the periodic law.
- RAMSAY, COLLIE, and Travers. *Compt. rend.*, **123**, pp. 214, 542; 1896; *Proc. Roy. Soc., Lond.*, **60**, p. 206; 1896; *Proc. Roy. Soc., Lond.*, **62**, p. 316; 1898: Attempt to separate helium into 2 or more components.
- RAMSAY, WM. *Rend. soc. chim. ital.* (2), **5**, pp. 137-150; 1907; *Chem. Abs.*, **7**, p. 3712: The rôle of helium in nature.
- SCHAUM, K. *Sitz.-Ber. Ges. Natw.*, pp. 43-44; 1904: The condition of helium in the sun.
- KEESOM, W. H. *Handl. Ned. Nat. Geneesk. Congres*, **11**, pp. 169-174; 1907; *Leiden Comm. Phys. Lab. Suppl.* 18 to 97-108, pp. 9-16; 1908: The properties of helium mixtures.
- COOKE, W. T. *Proc. Roy. Soc., Lond.*, **77**, pp. 148-155; 1906; *Z. phys. Chem.*, **55**, pp. 537-546; 1906: Experiments on the chemical behavior of argon and helium. Vapor densities of various elements in these gases at 1,200-1,300°.
- FORCRAND. *Compt. rend.*, **156**, pp. 1648-1651; 1913: The molecular heat of vaporization of substances boiling at low temperatures.
- HERTZ, G. *Ber. physik. Ges.*, **19**, pp. 268-288; 1917; *J. Chem. Soc.*, **114**, II, pp. 105-106; 1918; *Chem. Abs.*, **12**, p. 1723: Exchange of energy in the collisions between slowly moving electrons and molecules of gases. Calculated for helium and hydrogen which differ by a factor of 200.
- JEANS. *Phil. Mag.* (6), **8**, p. 692; 1904: The diameter of a helium molecule.
- RANKINE, A. O. *Phil. Mag.* (6), **40**, pp. 516-519; 1920; *Chem. Abs.*, **15**, p. 462.
- BROUGHALL, L. S. C. *Phil. Mag.* (6), **41**, pp. 872-876; 1921; *Chem. Abs.*, **15**, p. 2775: Some dimensions of the atom.
- DEWAR, J. *Engineering*, **87**, pp. 825-826; 1909; *Chem. abs.*, **4**, p. 5: Problems connected with helium. Principally the use of helium for producing low temperatures.
- DEWAR, J. *Rev. Sci.*, **49**, pp. 450-457; 1911: Experiments with helium at low temperatures and low pressures.
- EDITORIAL. *Engineering*, **99**, pp. 144-145; 1915; *Chem. Abs.*, **9**, p. 1269: Hydrogen and the rare gases. Review of lecture by J. Dewar.
- MOORE, R. B. *J. Frank. Inst.*, **191**, pp. 145-197; 1921: Helium, its history, properties, and commercial development.
- MOORE, R. B. *Bur. of Mines, Serial No.* 2363; 1922: Report of helium investigations of the Bureau of Mines.
- MOORE, R. B. *Natural Gas*, **111**, p. 14, 1922: Helium, its history, properties, and commercial development.
- MOORE, R. B. *The Carolina Chemist*, **8**, p. 7, 1922: Helium.
- LOCKYER, J. N. *The sun's place in nature*; 300 pp. The MacMillan Co., London, 1897.
- RAMSAY, W., and RUDORF, G. *Die Edelgase* (the noble gases) (book), Vol. II of Ostwald and Drucker's *Handbuch der allgemeinen chemie*, Leipzig Akad. Verlagsgesellschaft; *Chem. Abs.*, **12**, pp. 2482; 1918.
- RAMSAY, W. *Die Edlen und die Radioactiven Gase* (the noble and the radioactive gases) (book), Leipzig Akad. Verlagsgesellschaft; *Chem. Abs.*, **3**, p. 747.
- MOORE, R. B. *J. Franklin Inst.*, **186**, pp. 29-55; 1918; *Chem. Abs.*, **12**, p. 1607: Sir William Ramsay. An account of his contributions to chemistry and a bibliography of his papers.
- STREUDEL, V. *On Argon and Helium*. O. Riecker, Tübingen; 1901.



